


Coronavirus: A possible cause of reduced male fertility

Chuan Huang^{1,2} | Xiren Ji¹ | Wenjun Zhou¹ | Zhenghui Huang^{1,2} | Xiangjie Peng² | Liqing Fan^{1,2} | Ge Lin^{1,2} | Wenbing Zhu^{1,2} 

¹Institute of Reproductive and Stem Cell Engineering, Basic Medicine College, Central South University, Changsha, China

²Reproductive and Genetic Hospital of CITIC-Xiangya, Changsha, China

Correspondence

Wenbing Zhu, PhD, Reproductive & Genetic Hospital of CITIC-Xiangya. No. 87, Xiangya Road, Changsha, Hunan 410008, China.
Email: zhuwenbing0971@sina.com

Funding information

China Postdoctoral Science Foundation, Grant/Award Number: 2019M661521

Abstract

In lately December 2019, a novel coronavirus (SARS-CoV-2) outbreak occurred in Wuhan, PR China. It is a high contagious virus that has threatened human health worldwide. SARS-CoV-2 infection, termed COVID-19, causes rapidly developing lung lesions that can lead to multiple organ failure in a short period. Whenever a novel virus emerges, reproductive risk assessments should be performed after infection. In this review, we show that male fertility might be damaged by coronavirus associated with (i) direct cytopathic effects derived from viral replication and viral dissemination in the testis; and (ii) indirect damage to male fertility derived from immunopathology. In this review, we briefly describe the impaired fertility of humans and animals infected with coronaviruses to deduce the impact of the new coronavirus on male fertility. Together with information related to other coronaviruses, we extrapolate this knowledge to the new coronavirus SARS-CoV-2, which may have a significant impact on our understanding of the pathophysiology of this new virus.

KEYWORDS

coronavirus, fertility, male, SARS-CoV-2, testis

1 | INTRODUCTION

Coronaviruses are the largest family of positive-stranded RNA viruses, which includes 30 members at present. They are widely distributed in nature, including infections of humans and other mammals. In recent years, new coronaviruses have caused problems worldwide in cycles, such as severe acute respiratory syndrome coronavirus (SARS-CoV) occurring in 2002, and Middle East respiratory syndrome coronavirus (MERS-CoV) being first identified in 2012. In 2019, a new highly contagious virus broke out in Wuhan, Hubei province, China, termed SARS-CoV-2, representing the seventh member of enveloped RNA coronaviruses.¹ The 2019 novel coronavirus disease (COVID-19) caused by SARS-CoV-2 has common clinical manifestations such as fever, dry cough, and in severe cases, multiple organ damage.²⁻⁴

Regarding the critical molecule for SARS-CoV-2 transmission, the receptor angiotensin I-converting enzyme 2 (ACE2) for virus cell entry and transmembrane serine protease 2 TMPRSS2 for priming the S protein⁵ are co-expressed in the testis and male genital tract,⁶ which suggests a high possibility that the virus targets the testis and male genital tract during infection. It was reported that over 25 viruses could enter human semen and negatively affect spermatozoa or male fertility,⁷ such as HSV⁸ and HIV.⁹ Whether SARS-CoV-2 may have the same the effect on males is an important question that was not answered unambiguously in a preliminary investigation.¹⁰

To date, studies¹¹⁻¹³ have confirmed the absence of SARS-CoV-2 RNA in the semen of patients with COVID-19. Conversely, The results of Li et al's study were inconsistent with those of previous studies and detected of SARS-CoV-2 in 6 of 38 semen samples.¹⁴ Similarly, Yang et al reported that one case (1/12) with a high viral load was positive

Chuan Huang and Xiren Ji contributed equally to this work.

© 2020 American Society of Andrology and European Academy of Andrology

for viral RNA after post-mortem examinations of testicular tissue,¹⁵ which supported the idea that high viral loads in patients with severe disease symptoms might reach the threshold to cross the blood-testis barrier.¹⁶ On the other hand, a study showed that compared with patients with mild disease, patients with severe COVID-19 have significantly lower testosterone levels,¹⁷ suggesting that the co-expression of ACE2 and TMPRSS2 on Leydig cells might make them susceptible to SARS-CoV-2 infection and thus compromise testosterone secretion.¹⁸ However, considering the high false-negative results for SARS-CoV-2 using RT-PCR,¹⁹ as well as the limitation of the small sample size and selection bias mostly obtained from recovering mild cases,¹⁰ we still need to be cautious when evaluating these data. Nevertheless, it is well-known that coronaviruses can contribute to high morbidity and mortality in both humans and animals.^{20,21} A study has demonstrated orchitis in patients with SARS, with detrimental effects in the testis, suggesting that coronavirus can infect the male reproductive tract and impair male reproduction.²² SARS-CoV-2 and SARS-CoV share some common clinical manifestations, which supports the hypothesis that the new coronavirus might directly infect the testes and male reproductive system. Therefore, we should be vigilant about the impact on male reproduction in patients with COVID-19. In addition, the blood-testis barrier might allow the testes to act as a special reservoir to protect viruses against antiviral agents,²³ which is a key reason for considering the testes as a particularly important organ for study in the context of the SARS-CoV-2 pandemic, and it is especially important because the coronavirus family has been identified the culprit causing orchitis in both humans (SARS-CoV)²² and animals (feline coronavirus and avian coronavirus).^{24,25} Using evidence from previous studies of coronavirus-infected animals and humans, the implications of this review may help us to understand the impact of SARS-CoV-2 on male reproductive capacity.

2 | DIRECT VIRUS-INDUCED CYTOPATHIC EFFECTS

Various viruses can replicate in the male reproductive tract, such as HEV²⁶ and ZIKV,^{27,28} which eventually lead to testicular atrophy and male infertility. Viral infection of the male genital tract can provide insights into possible male fertility impairment after SARS-CoV-2 infection. SARS-CoV-2 enters cells by binding ACE2 and via priming by TMPRSS2. ACE2 is a membrane-associated secretase that is expressed primarily on endothelial cells and is the host cell receptor for SARS and SARS-CoV-2.²⁹⁻³¹ Notably, ACE2 is highly tissue-specific, with significant levels being detected only in the heart, kidneys, testes, and gastrointestinal tract.³²⁻³⁴ In the testes, ACE2 is expressed only in spermatogenic cells and testis somatic cells, suggesting a high potential for testicular damage and spermatogenesis disruption when the virus combines with this metalloprotease.³⁵ TMPRSS2, as an essential protease for viral infection, is highly expressed in spermatogonia and spermatids.¹⁸ The co-expression of ACE2 and TMPRSS2 in spermatogonia and Leydig cells implied that the testis might be a high-risk organ that is vulnerable to SARS-CoV-2 infection, which

might result in testicular degeneration and male infertility. SARS coronaviruses, whose expressed proteins share 76% amino acid sequence identity with those of SARS-CoV-2, were detected in testis somatic cells.³⁶ This observation supports the hypothesis that the SARS-CoV-2 might concentrate on testis cells to dysregulate their function.

2.1 | Direct virus-induced damage of the testis

Viral replication in cells contributes directly to microscopy-detected lesions, which eventually result in spermatogonia necrosis,²⁶ such as in a ram model of infection by Bluetongue virus (BTV) (an arbovirus of ruminants), which showed testicular parenchyma damage and the destruction of the Sertoli cells caused by viral replication-induced cytopathic effects.³⁷ Coronaviruses might use a similar mechanism in humans to impair male fertility. ACE2 is the crucial determinant of coronavirus infection, tissue tropism, and subsequent viral replication.^{38,39} The expression pattern of ACE2 in adult human testis at the level of single-cell transcriptomes was shown to be predominantly enriched in Leydig and Sertoli cells.⁶ Besides, alternative receptor Basigin (BSG) and protease Cathepsin L (CTSL) were also detected in Leydig cells,⁴⁰ which can mediate SARS-CoV-2 into cells. Data from autopsies of 12 patients with COVID-19 showed a dramatic reduction in Leydig cells in the interstitium,¹⁵ supporting the speculation that SARS-CoV-2 could display tropism for Leydig cells, ultimately leading to ultrastructural lesions and decreased numbers of Leydig cells. Leydig cells occur in clusters between blood vessels and seminiferous tubules, producing the vast majority of androgens in men.⁴¹ The replication of SARS-CoV-2 in testosterone-producing Leydig cells might disrupt testosterone production. Indeed, a recent study confirmed that patients with COVID-19 suffered hypogonadotropic hypogonadism as the disease progressed, implying that the secretory function of Leydig cells might be impaired by the novel coronavirus.¹⁷ Testosterone is essential to preserve male fertility and to support Sertoli cell maturation and the development of Leydig cells.⁴² Extensive evidence from clinical and laboratory studies implied that testosterone deficiency is accompanied by atrophy of the testicular parenchyma and degradation in the seminiferous tubules,^{43,44} and in summary, testosterone is necessary for men to maintain the blood-testis barrier, spermatogenesis, and fertility. Alterations in male sex hormone levels induced by SARS-CoV-2 might negatively affect male reproduction. Therefore, special attention should be paid to andrology examinations and hormone assessments on men recovering from COVID-19, as well as exploring the possible long-term outcomes of SARS-CoV-2 infection.

Sertoli cells are the only somatic cells in the tubules that directly contact with spermatogenic cells and control the differentiation of spermatogenic cells via paracrine signals.⁴⁵ Inhibin B is secreted by Sertoli cells, and compared with follicle-stimulating hormone (FSH) or luteinizing hormone (LH), it is an ideal marker for spermatogenesis and a better indicator of sterility.^{46,47} SARS-CoV-2 has a high affinity for human ACE2, which suggests that the virus might concentrate on Sertoli cells. Indeed, inhibin B levels decreased after hepatitis E

virus infection in mice, which was attributed to damage of the Sertoli cells in the testes.²⁶ Accumulating evidence suggests that the coronavirus family has an affinity for these testes cells, for example, avian infectious bronchitis virus (IBV), a subtype of coronavirus, can cause acute respiratory infections in birds,⁴⁸ and was detected in Sertoli cells of the testes of infected roosters using immunofluorescence.⁴⁹ Roosters vaccinated with live attenuated IBV showed significantly reduced serum androgen concentrations compared with non-vaccinated roosters and could cause infertility in roosters.⁵⁰ Considering that IBV causes a similar severe acute respiratory syndrome to SARS-CoV-2, we hypothesized that the same mechanism might be used by SARS-CoV-2 to spread in Sertoli cells. In addition, roosters vaccinated with live attenuated IBV might provide an animal model of how SARS-CoV-2 replicates in cells and causes pathogenic effects in the testis.

Additionally, coronavirus might directly disrupt the microenvironment of the testis that supports spermatogenesis. In CoV-infected roosters, histological analysis revealed the disruption of seminiferous tubules and loss of the basement membrane, leading to the destruction of the spermatogenesis microenvironment, which contributed to the reduction of the live sperm concentration.⁵¹ Thus, testicular degeneration is possibly the result of several overlapping factors once a coronavirus infects the male genital tract, and this might also be the case for SARS-CoV-2.

2.2 | Virus-induced damage of spermatogenesis directly

Viruses can be detected in semen directly. SARS-CoV-2 RNA has been isolated from rectal swabs and respiratory tract swabs.⁵² Currently, the question of whether the virus can infect semen needs an answer. According to a recent study of scRNA-seq data in adult human testes, ACE2 and TMPRSS2 are highly co-expressed in spermatogonia, which are enriched in the gene ontology (GO) categories relating to viral reproduction and transmission.⁶ Therefore, it is reasonable to hypothesize that there is a high risk of SARS-CoV-2 presence in seminal fluid.⁵³ However, a few case reports have investigated this issue, and the presence of SARS-CoV-2 in semen remains ambiguous.¹¹⁻¹⁴ Notably, gene ontology (GO) enrichment analysis illustrated that cell junction and immunity-related GO terms were enriched in ACE2-positive Leydig/Sertoli cells; therefore, cell-cell junctions might allow the transfer of SARS-CoV-2,⁶ which might represent one explanation of the highly contagious nature of this novel coronavirus and could have implications for sexual and reproductive behavior.⁵⁴ Taken together, there is a critical need to verify virus infection semen and whether sexual transmission of SARS-CoV-2 can indeed occur.

With regard to research on coronavirus infection animals, IBV has been isolated from testicles and semen in roosters,^{51,55} and when insemination using IBV-spiked semen was performed, IBV RNA could be detected in all the hens, and the weight of eggs laid by the hens inseminated with IBV-spiked semen was significantly reduced.⁵⁵ Knowledge of other coronaviruses present in semen might encourage researchers to look at semen and sexual transmission,

to determine whether SARS-CoV-2 can be sexually transmitted like IBV; however, the results will need to be cautiously interpreted. Nevertheless, we should remain vigilant to this possibility, which has important implications in reproductive medicine, especially viral transmission facilitated by ART, such as intracytoplasmic sperm injection (ICSI),⁵⁶ sperm cryopreservation, and the prevention of transmission. During this epidemic, sperm cryobank must introduce precautionary measures: First, we recommend that semen from SARS-CoV-2-positive men is cryopreserved in a highly secure, separate container, such as a vapor cryostorage tank. Secondly, all donors must undergo mandatory SARS-CoV-2 testing. Thirdly, abstinence or condom use might be considered as preventive measure for patients with COVID-19.

Virus binding to ACE2-expressing spermatogonia would disrupt spermatogenesis.⁶ Rooster vaccination with coronavirus caused a significant reduction in daily sperm production.^{25,57} A recent report also confirmed that semen quality parameters were impaired in patients with moderate infection of COVID-19.⁵⁸ Hence, the risk of SARS-CoV-2 virus infection to semen parameters may not be negligible. Notably, the long-term impact on semen parameters of SARS-CoV-2 infection, as well as semen examination, is required during follow-up patients recovering from COVID-19, especially men who plan to have children.

2.3 | Direct virus-induced damage of the epididymis

In animal models of coronavirus infection, one of the major clinical symptoms is epididymal stone formation.^{51,59,60} IBV replication in roosters' testes might result in severe cellular micropathological damage, which in the long-term can lead to the presence of epididymal stones. The presence of stones is associated with reduced fertility and adverse effects on sperm function,⁶¹ eventually resulting in the collapse of the seminiferous tubules and cessation of spermatogenesis. The epididymis is a crucial region for sperm maturation, which is pivotal for spermatozoa to obtain the motile ability and fertile capacity. Dysfunction in this area can compromise sperm maturation and further impair sperm quality, such as decreased sperm motility, increased DNA damage, changed membrane lipids, and the acrosome reaction.⁶² If the behavior of coronavirus infection in humans is similar to that in animal models, we should pay attention to the epididymis to protect it from SARS-CoV-2-induced destruction.

In view of these results, we suggest prompting a comprehensive genitourinary examination for patients with COVID-19, including alterations in semen parameters, such as the acrosome reaction, DNA damage, and sperm motility.

3 | INDIRECT IMMUNE-MEDIATED DAMAGE TO MALE FERTILITY

The testicle is an immunologically privileged organ, the blood-testis barrier (BTB) protects the testes against pathogen invasion.⁶³ In

healthy fertile men, various immune cells and cytokines produced by non-immune cells are indispensable to ensure male fertility,⁶⁴ in which they maintain the testicular microenvironment balance and male reproductive health within the intricate and active environment of the seminiferous epithelium. Testicular tissue development benefits from immune cells and their cytokines, and the immune response is critical to control and eliminate viral infection.⁶⁵ Cytokines are important for the immune response to viral infections by regulating the expansion and location of leukocytes. However, infection and inflammation might disrupt the immune balance in the body, either through immune insufficiency or overactivation, possibly leading to devastating effects in humans.⁶⁶ Immune pathology associated with an uncontrolled immune response might give rise to testicular parenchyma destruction when the BTB is damaged by virus infection,⁶⁷ and any associated functional impairment could lead to male infertility.

3.1 | Cytokine-mediated infertility

SARS-CoV-2 has proven effects on multiple organs throughout the body,⁶⁸ accompanied by immunopathological reactions and high cytokine storms. In the plasma of patients with COVID-19 in intensive care units, higher plasma levels of cytokines were detected, implying that a cytokine storm might aggravate the infection in patients with COVID-19.^{2,3} Actually, this coincided with the research that patients with COVID-19 presented a typical profile of hyperinflammation, such as TNF- α , IL-6, and IL-1 β .⁶⁹ Cytokines are beneficial to testicular function and sperm production, as well as testicular immunity privilege.⁷⁰⁻⁷² However, a high concentration of inflammatory cytokines could contribute to the progression of sexual dysfunction.⁷³ Thus, a change in cytokine production can lead to fertility problems.^{66,74} Cytokine-mediated suppression of the hypothalamic-pituitary-testicular axis could lead to a decrease in serum testosterone, such as IL1 leading to inactivation of the P450/c17 lyase that converts progestins into androgens in immunopathogenesis, which will result in decreased testosterone and sperm production.^{60,75-77} This corroborated the results showing a dramatic decline serum testosterone in 17 patients with severe COVID-19, which might even predict poor progression of COVID-19 infection.⁷⁸ With a history of COVID-19 disease, SARS-CoV-2 infection can attribute to male hypogonadism,⁷⁹ thus it is recommended to measure testosterone levels when a patient is detected as positive for SARS-CoV-2 RNA and conduct appropriate testosterone treatment if necessary. Studies detected dramatic increases in IL6 levels in patients with COVID-19.^{80,81} Immunopathologically, high IL6 expression correlates with a systemic inflammatory milieu that disrupts the integrity of the blood-testis barrier.⁸² As a result of blood-testis dissemination, the virus might damage testicular tissue directly. Furthermore, COVID-19-induced changes to the cytokine microenvironment might even lead to testicular cancer,⁶⁴ which could have long-term adverse effects on the recovery of patients, and represents a second long-term matter of concern. Hence, it should be noted that the cytokine

storm introduced by SARS-CoV-2 could be associated with immunopathogenesis, which might contribute to testicular dysfunction and reduced male fertility. Nevertheless, this hypothesis requires follow-up confirmation, and the exploration of possible short- and long-term consequences on their andrological health.

3.2 | Inflammation-mediated infertility

The blood-testicular barrier might not be a perfect barrier to viruses under systemic or local inflammation.⁷ To eliminate the virus infection, an inflammatory cytokines storm can recruit leukocytes, resulting in inflammation characterized by leukocyte infiltration in the interstitial tissue of the testes, which, as a feature of human testicular orchitis, might lead to male infertility. Actually, there is a high risk of that men with SARS-CoV-2 might suffer from an orchitis-like syndrome.³⁵ Pan et al confirmed that six patients (19%) with COVID-19 suffered from orchitis.¹¹ Recently, a study of 12 deceased patients with COVID-19 also revealed viral orchitis characteristics, with T lymphocyte intrusion into the testicular parenchyma, accompanied by significant seminiferous tubular injury.¹⁵ Interestingly, the histopathological features of the testes in patients with SARS also overlap with those in patients with COVID-19: all testes being full of leukocyte infiltration and wide-ranging germ cell deterioration, with thickened basement membranes,⁸³ which supports the hypothesis that the coronavirus-induced adaptive immune response might play a vital role in the course of testicular damage and eventually affect fertility. Theoretically, attributed to the hypercoagulable state of vasculitis in patients with COVID-19, the testicular damage could be result of testicular segmental vascularization.⁸⁴ One study has shown evidence of direct SARS-CoV-2 infection of endothelial cells and diffuse endothelial inflammation,⁸⁵ Endothelial dysfunction may be subsequent to organ ischemia,⁸⁶ which might provide a rationale for one study that described ischemia-related priapism in a patient with COVID-19,⁸⁷ suggesting that vasculitis-orchitis might have a crucial role in the development of the testicular injury caused by SARS-CoV-2 infection. Moreover, the intrusion of CD68 + macrophages into the interstitial tissue of the testes could contribute to a decline in steroidogenesis and testosterone,⁶⁶ and the change in the hormonal profile might contribute to susceptibility to SARS-CoV-2 infection, leading to a more profound pathophysiological role in COVID-19 patients.⁸⁸

During the SARS-CoV-2 outbreak, SARS-CoV-2-infected cats also presented a profile of testicular atrophy⁸⁹ and were reported to have acquired the infection from humans.⁹⁰ Furthermore, studies on chickens infected with coronavirus IBV also showed that immune cells infiltrated into the interstitium of the testis, which was responsible for the reduced fertility.^{25,91}

In summary, the coronavirus-induced adaptive immune response might lead to testicular damage and endocrine abnormality, eventually disrupting spermatogenesis in patients recovering from COVID-19. However, this hypothesis remains to be confirmed and studies should be undertaken to establish an animal model to determine the underlying pathophysiological mechanisms and

to mitigate the risk of testicular injury during COVID-19 disease. Precautions against SARS-CoV-2-induced male infertility should be taken.

3.3 | Antibody-mediated infertility by SARS-CoV-2

In SARS-infected testes, Immunohistochemistry analysis showed a large amount of IgG precipitation in the seminiferous epithelium of the testis, as well as in degraded germ cells and Sertoli cells, suggesting that the extensive IgG triggered by a secondary autoimmune response might aggravate the testicular damage.²² In addition, deposits of IgG are associated with autoimmune orchitis (EAO),⁹² which might activate immune cells in the host to produce antibodies against the virus, as well as introducing antibodies into semen.⁹³ In patients with COVID-19, the positive rate of IgG reached 100%,⁹⁴ especially antiphospholipid antibodies,⁹⁵ which are antisperm antibodies that could interfere with fertilization,⁹⁶ suggesting that male patients with COVID-19 should be cautioned against the adverse effects of a high IgG titer on their reproduction ability.

In healthy testis tissue, immune cells and cytokines are beneficial for the development of spermatogonia. However, the immune imbalance associated with infection and inflammation can contribute to male sterility. Overall, in addition to the pathogenic effects of coronavirus, the host-induced immune response against the virus also plays an important role in the overall disease process.

3.4 | High fever and steroid-mediated infertility

It is generally believed that high fever can be detrimental to the normal function of the testes. Fever is one of the notable features of COVID-19³ and thus might play an important role in testicular dysfunction. Germ cells can develop at a normal pace at temperatures less than 37.8°C; however, higher temperatures might cause irreversible damage to germ cells. Research confirms that high temperatures can lead to the meiotic arrest of germ cells.⁹⁷ In general, increased body temperature has a negative influence on spermatogenesis and may ultimately lead to male infertility.

In addition, patients with COVID-19 were almost all affected by SARS-CoV-2-related stress and were advised to use steroids (methylprednisolone, 1-2 mg/kg per day)² to treat SARS-CoV-2; however, the stress⁹⁸ and corticosteroid therapy might have adverse effects on sexual function, such as reduced libido and erectile dysfunction.^{99,100} Leydig cells were also proven to be dysfunctional in glucocorticoid-treated rats.¹⁰¹ Therefore, these observations suggest that the assessment of fertility in patients with COVID-19 is imperative.

4 | CONCLUSION

This review obtained clues from basic research on other viruses to understand how the novel SARS-CoV-2 virus might generate

pathogenic effects in male fertility. We highlighted that male fertility might be highly vulnerable to SARS-CoV-2 infection. Infection with this novel virus not only seriously threatens an individual's overall health, but also might lead to male infertility. Perspectives gained from multi-organ research during the recent epidemic raises the possibility that damage to the male reproductive tract might be an underappreciated result of SARS-CoV-2 infection. Therefore, more attention should be paid to the effects on male fertility of SARS-CoV-2 infection, and should this causal link between SARS-CoV-2 infection and male infertility be confirmed, male patients should consider cryopreserving their spermatozoa to preserve fertility.

ACKNOWLEDGEMENTS

This study was supported by a grant from China Postdoctoral Science Foundation (2019M661521).

CONFLICT OF INTEREST

None declared.

AUTHORS' CONTRIBUTION

Wenbing Zhu and Chuang Huang conceived and designed the study. Chuang Huang and Xiren Ji drafted the manuscript. Wenjun Zhou, Zhenghui Huang, Xiangjie Peng, Liqing Fan, and Ge Lin revised the drafts. All authors approved the final version of the manuscript.

ORCID

Wenbing Zhu  <https://orcid.org/0000-0002-9352-2785>

REFERENCES

- Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. 2020;382(8):727-733.
- Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507-513.
- Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*. 2020;395(10223):497-506.
- Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *New Engl J Med*. 2020;382(13):1199-1207.
- Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell*. 2020;181(2):271-280.
- Wang Z, Xu X. scRNA-seq profiling of human testes reveals the presence of the ACE2 receptor, a target for SARS-CoV-2 infection in Spermatogonia, Leydig and Sertoli Cells. *Cells*. 2020;9(4):920.
- Salam AP, Horby PW. The breadth of viruses in human semen. *Emerg Infect Dis*. 2017;23(11):pp. 1922-1924.a.
- Klimova RR, Chichev EV, Naumenko VA, et al. Herpes simplex virus and cytomegalovirus in male ejaculate: herpes simplex virus is more frequently encountered in idiopathic infertility and correlates with the reduction in sperm parameters. *Vopr Virusol*. 2010;55(1):27-31.
- Garrido N, Meseguer M, Remohi J, Simon C, Pellicer A. Semen characteristics in human immunodeficiency virus (HIV)- and hepatitis C (HCV)-seropositive males: predictors of the success of viral removal after sperm washing. *Hum Reprod*. 2005;20(4):1028-1034.
- Paoli D, Pallotti F, Turriziani O, et al. SARS-CoV-2 presence in seminal fluid: Myth or reality. *J Androl*. 2020.

11. Pan F, Xiao X, Guo J, et al. No evidence of severe acute respiratory syndrome-coronavirus 2 in semen of males recovering from coronavirus disease 2019. *Fertil Steril*. 2020;113(6):1135-1139.
12. Paoli D, Pallotti F, Colangelo S, et al. Study of SARS-CoV-2 in semen and urine samples of a volunteer with positive naso-pharyngeal swab. *J Endocrinol Invest*. 2020;1-4.
13. Song C, Wang Y, Li W, et al. absence of 2019 novel coronavirus in semen and testes of COVID-19 patients. *Biol Reprod*. 2020;103(1):4-6.
14. Li D, Jin M, Bao P, Zhao W, Zhang S. Clinical characteristics and results of semen tests among men with coronavirus disease 2019. *JAMA Netw Open*. 2020;3(5):e208292.
15. Yang M, Chen S, Huang B, et al. Pathological findings in the testes of COVID-19 patients: clinical implications. *Europ Urol Focus*. 2020;6(5):1124-1129.
16. Liu Y, Yan LM, Wan L, et al. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis*. 2020;20(6):656-657.
17. Rastrelli G, Di Stasi V, Inglese F, et al. Low testosterone levels predict clinical adverse outcomes in SARS-CoV-2 pneumonia patients. *Andrology*. 2020;1-11.
18. Wang Z, Xu X. scRNA-seq Profiling of human testes reveals the presence of the ACE2 receptor, a target for SARS-CoV-2 infection in Spermatogonia, leydig and sertoli cells. *Cells*. 2020;9(4):920.
19. Kelly JC, Dombrowski M, O'Neil-Callahan M, Kernberg AS, Frolova AI, Stout MJ. False-negative COVID-19 testing: considerations in obstetrical care. *Am J Obstet Gynecol MFM*. 2020;2(3):100130.
20. Li W, Shi Z, Yu M, et al. Bats are natural reservoirs of SARS-like coronaviruses. *Science*. 2005;310(5748):676-679.
21. Kupferschmidt K. Emerging diseases. Researchers scramble to understand camel connection to MERS. *Science*. 2013;341(6147):702.
22. Xu J, Qi L, Chi X, et al. Orchitis: a complication of severe acute respiratory syndrome (SARS). *Biol Reprod*. 2006;74(2):410-416.
23. Shastri A, Wheat J, Agrawal S, et al. Delayed clearance of SARS-CoV2 in male compared to female patients: High ACE2 expression in testes suggests possible existence of gender-specific viral reservoirs. *MedRxiv*. 2020.
24. Sigurðardóttir ÓG, Kolbjørnsen Ø, Lutz H. Orchitis in a cat associated with coronavirus infection. *J Comp Pathol*. 2001;124(2-3):219-222.
25. Boltz DA, Zimmerman CR, Nakai M, Bunick D, Scherba G, Bahr JMAd. Epididymal stone formation and decreased sperm production in roosters vaccinated with a killed strain of avian infectious bronchitis virus. *Avian Dis*. 2006;50(4):594-598.
26. Situ J, Wang W, Long F, et al. Hepatitis E viral infection causes testicular damage in mice. *Virology*. 2020;541:150-159.
27. Govero J, Esakky P, Scheaffer SM, et al. Zika virus infection damages the testes in mice. *Nature*. 2016;540(7633):438-442.
28. Ma W, Li S, Ma S, et al. Zika virus causes testis damage and leads to male infertility in mice. *Cell*. 2017;168(3):542.
29. Li W, Moore MJ, Vasilieva N, et al. Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature*. 2003;426(6965):450-454.
30. Hoffmann M, Kleine-Weber H, Krüger N, Müller M, Drosten C, Pöhlmann S. The novel coronavirus 2019 (2019-nCoV) uses the SARS-coronavirus receptor ACE2 and the cellular protease TMPRSS2 for entry into target cells. *bioRxiv*. 2020:2020.2001.2031.929042.
31. Zhou P, Yang X-L, Wang X-G, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020;579(7798):270-273.
32. Tipnis SR, Hooper NM, Hyde R, Karran E, Christie G, Turner AJ. A human homolog of angiotensin -converting enzyme cloning and functional expression as a captopril-insensitive carboxypeptidase. *J Biol Chem*. 2000;275(43):33238-33243.
33. Harmer D, Gilbert M, Borman R, Clark KL. Quantitative mRNA expression profiling of ACE 2, a novel homologue of angiotensin converting enzyme. *FEBS Lett*. 2002;532(1):107-110.
34. Douglas GC, O'Bryan MK, Hedger MP, et al. The novel angiotensin-converting enzyme (ACE) homolog, ACE2, is selectively expressed by adult Leydig cells of the testis. *Endocrinology*. 2004;145(10):4703-4711.
35. Corona G, Baldi E, Isidori AM, et al. SARS-CoV-2 infection, male fertility and sperm cryopreservation: a position statement of the Italian Society of Andrology and Sexual Medicine (SIAMS) (Societa Italiana di Andrologia e Medicina della Sessualita). *J Endocrinol Invest*. 2020;43(8):1153-1157.
36. Zhao J-M, Zhou G-D, Sun Y-I, et al. Clinical pathology and pathogenesis of severe acute respiratory syndrome. *Chinese J Clin Hepatology*. 2003;17(3):217-221.
37. Puggioni G, Pintus D, Melzi E, et al. Testicular degeneration and infertility following arbovirus infection. *J Virol*. 2018;92(19):e01131-18.
38. Shang J, Ye G, Shi K, et al. Structural basis of receptor recognition by SARS-CoV-2. *Nature*. 2020;581(7807):221-224.
39. Perlman S, Netland J. Coronaviruses post-SARS: update on replication and pathogenesis. *Nat Rev Microbiol*. 2009;7(6):439-450.
40. Stanley KE, Thomas E, Leaver M, Wells D. Coronavirus disease-19 and fertility: viral host entry protein expression in male and female reproductive tissues. *Fertil Steril*. 2020;114(1):33-43.
41. Neto FT, Bach PV, Najari BB, Li PS, Goldstein M. Spermatogenesis in humans and its affecting factors. *Semin Cell Dev Biol*. 2016;59:10-26.
42. Guan X, Chen F, Chen P, et al. Effects of spermatogenic cycle on Stem Leydig cell proliferation and differentiation. *Mol Cell Endocrinol*. 2019;481:35-43.
43. Eroschenko VP, Wilson WO, Siopes TD. Function and histology of testes from aged coturnix maintained on different photoperiods. *J Gerontol*. 1977;32(3):279-285.
44. Uraki R, Hwang J, Jurado KA, et al. Zika virus causes testicular atrophy. *Sci Advanc*. 2017;3(2):1602899.
45. Chen S-R, Liu Y-X. Regulation of spermatogonial stem cell self-renewal and spermatocyte meiosis by Sertoli cell signaling. *Reproduction*. 2015;149(4).
46. Anderson RA, Irvine DS, Balfour C, Groome NP, Riley SC. Inhibin B in seminal plasma: testicular origin and relationship to spermatogenesis. *Hum Reprod*. 1998;13(4):920-926.
47. Mahmoud AM, Comhaire FH, Depuydt CE. The clinical and biologic significance of serum inhibins in subfertile men. *Reprod Toxicol*. 1998;12(6):591-599.
48. Cavanagh D. Nidovirales: a new order comprising Coronaviridae and Arteriviridae. *Arch Virol*. 1997;142(3):629-633.
49. Gallardo RA, Hoerr FJ, Berry WD, van Santen VL, Toro H. Infectious bronchitis virus in testicles and venereal transmission. *Avian Dis*. 2011;55(2):255-258.
50. Europe JRC. history, current situation and control measures for infectious bronchitis. *Brazilian J Poultry Sci*. 2010;12(2):125-128.
51. Villarreal L, Brandão PE, Chacón JL, et al. Orchitis in roosters with reduced fertility associated with avian infectious bronchitis virus and avian metapneumovirus infections. *Avian Dis*. 2007;51(4):900-904.
52. Guan W-J, Ni Z-Y, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *New Engl J Med*. 2020;382(18):1708-1720.
53. Verma S, Saksena S, Sadri-Ardekani H. ACE2 receptor expression in testes: implications in coronavirus disease 2019 pathogenesis. *Biol Reprod*. 2020;103:449-451.
54. Aversa A, Jannini EA. COVID-19, or the triumph of monogamy? *Minerva Endocrinol*. 2020;45(2):77-78.

55. Gallardo RA, Hoerr FJ, Berry WD, Santen VL, Toro H. Infectious bronchitis virus in testicles and venereal transmission. *Avian Dis*. 2011;55(2):255-258.
56. Perry MJ, Arrington S, Neumann LM, Carrell D, Mores CN. It is currently unknown whether SARS-CoV-2 is viable in semen or whether COVID-19 damages spermatozoa. *Andrology*. 2020.
57. Boltz DA, Nakai M, Bahr JM. Avian infectious bronchitis virus: a possible cause of reduced fertility in the rooster. *Avian Dis*. 2004;48(4):909-915.
58. Holtmann N, Edimiris P, Andree M, et al. Assessment of SARS-CoV-2 in human semen—a cohort study. *Fertil Steril*. 2020;114(2):233-238.
59. Benyeda Z, Mató T, Süveges T, et al. Comparison of the pathogenicity of QX-like, M41 and 793/B infectious bronchitis strains from different pathological conditions. *Avian Pathol*. 2009;38(6):449-456.
60. Boltz DA, Nakai M, Bahr JM. Avian infectious bronchitis virus: a possible cause of reduced fertility in the rooster. *Avian Dis*. 2004;48(4):909-915.
61. Villarreal LYB, Brandão PE, Chacón JLV, et al. Orchitis in roosters with reduced fertility associated with avian infectious bronchitis virus and avian metapneumovirus infections. *Avian Dis*. 2007;51(4):900-904.
62. Liu Y, Ding Z. Obesity, a serious etiologic factor for male subfertility in modern society. *Reproduction*. 2017;154(4).
63. Li N, Wang T, Han D. Structural, cellular and molecular aspects of immune privilege in the testis. *Front Immunol*. 2012;3:152.
64. Loveland KL, Klein B, Pueschl D, et al. Cytokines in male fertility and reproductive pathologies: immunoregulation and beyond. *Front Endocrinol*. 2017;8:307.
65. Zhao S, Zhu W, Xue S, Han D. Testicular defense systems: immune privilege and innate immunity. *Cell Mol Immunol*. 2014;11(5):428-437.
66. Hedger MP, Meinhardt A. Cytokines and the immune-testicular axis. *J Reprod Immunol*. 2003;58(1):1-26.
67. Mruk DD, Cheng CY. The mammalian blood-testis barrier: its biology and regulation. *Endocr Rev*. 2015;36(5):564-591.
68. Liu Q, Wang RS, Qu GQ, et al. Gross examination report of a COVID-19 death autopsy. *Fa Yi Xue Za Zhi*. 2020;36(1):21-23.
69. Pedersen SF, Ho Y-C. SARS-CoV-2: a storm is raging. *J Clin Invest*. 2020;130(5):2202-2205.
70. Lui W-Y, Lee WM, Cheng CY. Transforming growth factor- β 3 perturbs the inter-sertoli tight junction permeability barrier in vitro possibly mediated via its effects on occludin, zonula occludens-1, and claudin-11. *Endocrinology*. 2001;142(5):1865-1877.
71. Dobashi M, Fujisawa M, Yamazaki T, Okada H, Kamidono S. Distribution of intracellular and extracellular expression of transforming growth factor-beta1 (TGF-beta1) in human testis and their association with spermatogenesis. *Asian J Androl*. 2002;4(2):105-109.
72. Itman C, Mendis SHS, Barakat BM, Loveland KAL. All in the family: TGF- β family action in testis development. *Reproduction*. 2006;132(2):233-246.
73. Maiorino MI, Bellastella G, Giugliano D, Esposito K. From inflammation to sexual dysfunctions: a journey through diabetes, obesity, and metabolic syndrome. *J Endocrinol Invest*. 2018;41(11):1249-1258.
74. Ochsenkühn R, O'Connor AE, Hirst JJ, Baker HWG, Kretser DMD, Hedger MP. The relationship between immunosuppressive activity and immunoregulatory cytokines in seminal plasma: Influence of sperm autoimmunity and seminal leukocytes. *J Reprod Immunol*. 2006;71(1):57-74.
75. Hales DB. Interleukin-1 inhibits Leydig cell steroidogenesis primarily by decreasing 17 α -hydroxylase/C17-20 lyase cytochrome P450 expression. *Endocrinology*. 1992;131(5):2165-2172.
76. Janssen SJ, Kirby JD, Hess RA, et al. Identification of epididymal stones in diverse rooster populations. *Poult Sci*. 2000;79(4):568-574.
77. Mahecha GAB, Oliveira CA, Balzuweit K, Hess RA. Epididymal lithiasis in roosters and efferent ductule and testicular damage. *Reproduction*. 2002;124(6):821-834.
78. Pozzilli P, Lenzi A. Commentary: Testosterone, a key hormone in the context of COVID-19 pandemic. *Metabolism*. 2020;108:154252.
79. Wambier CG, Goren A. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection is likely to be androgen mediated. *J Am Acad Dermatol*. 2020;83(1):308-309.
80. Wan S, Yi Q, Fan S, et al. Characteristics of lymphocyte subsets and cytokines in peripheral blood of 123 hospitalized patients with 2019 novel coronavirus pneumonia (NCP). *medRxiv*. 2020.
81. Liu J, Li S, Liu J, et al. Longitudinal characteristics of lymphocyte responses and cytokine profiles in the peripheral blood of SARS-CoV-2 infected patients. *Ebiomedicine*. 2020;55:102763.
82. Zhang H, Yin Y, Wang G, Liu Z, Liu L, Sun F. Interleukin-6 disrupts blood-testis barrier through inhibiting protein degradation or activating phosphorylated ERK in Sertoli cells. *Sci Rep*. 2015;4(1):4260.
83. Xu J, Qi L, Chi X, et al. Orchitis: a complication of severe acute respiratory syndrome (SARS). 2006; 74(2):410-416.
84. Leisman DE, Deutschman CS, Legrand M. Facing COVID-19 in the ICU: vascular dysfunction, thrombosis, and dysregulated inflammation. *Intens Care Med*. 2020;46(6):1105-1108.
85. Varga Z, Flammer AJ, Steiger P, et al. Endothelial cell infection and endotheliitis in COVID-19. *The Lancet*. 2020;395(10234):1417-1418.
86. Bonetti PO, Lerman LO, Lerman A. Endothelial dysfunction: a marker of atherosclerotic risk. *Arteriosclerosis Thromb Vasc Biol*. 2003;23(2):168-175.
87. Lamamri M, Chebbi A, Mamane J, et al. Priapism in a patient with coronavirus disease 2019 (COVID-19): a case report. *Am J Emerg Med*. 2020.
88. Salonia A, Corona G, Giwercman A, et al. SARS-CoV-2, testosterone and frailty in males (PROTEGGIMI): A multidimensional research project. *Andrology*. 2020.
89. Sigurdardottir OG, Kolbjornsen O, Lutz H. Orchitis in a cat associated with coronavirus infection. *J Comp Pathol*. 2001;124(2-3):219-222.
90. Zhang Q, Zhang H, Huang K, et al. SARS-CoV-2 neutralizing serum antibodies in cats: a serological investigation. *bioRxiv*. 2020.
91. Benyeda Z, Szeredi L, Mató T, et al. Comparative histopathology and immunohistochemistry of QX-like, Massachusetts and 793/B serotypes of infectious bronchitis virus infection in chickens. *J Comp Pathol*. 2010;143(4):276-283.
92. Itoh M, Hiramane C, Tokunaga Y, Mukasa A, Hojo K. A new murine model of autoimmune orchitis induced by immunization with viable syngeneic testicular germ cells alone. II. Immunohistochemical findings of fully-developed inflammatory lesion. *Autoimmunity*. 1991;10(2):89-97.
93. Moldoveanu Z, Huang WQ, Kulhavy R, Pate MS, Mestecky J. Human male genital tract secretions: both mucosal and systemic immune compartments contribute to the humoral immunity. *J Immunol*. 2005;175(6):4127-4136.
94. Long QX, Liu BZ, Deng HJ, et al. Antibody responses to SARS-CoV-2 in patients with COVID-19. *Nat Med*. 2020;26(6):845-848.
95. Connell NT, Battinelli EM, Connors JM. Coagulopathy of COVID-19 and antiphospholipid antibodies. *J Thromb Haemost*. 2020.
96. Chiu WWC, Chamley LW. Clinical associations and mechanisms of action of antisperm antibodies. *Fertil Steril*. 2004;82(3):529-535.
97. Xu J, Xu Z, Jiang Y, Qian X, Huang Y. Cryptorchidism induces mouse testicular germ cell apoptosis and changes in bcl-2 and bax protein expression. *J Environment Pathol Toxicol Oncol*. 2000;19:25-33.

98. Chan JC, Morgan CP, Leu NA, et al. Reproductive tract extracellular vesicles are sufficient to transmit intergenerational stress and program neurodevelopment. *Nat Commun*. 2020;11(1):1-13.
99. Ln C, Am M, Mm D, et al. Glucocorticoids: their role on gonadal function and LH secretion. *Minerva Endocrinol*. 1996;21(2):43-46.
100. Scaroni C, Favia G, Lumachi F, et al. Unilateral adrenal tumor, erectile dysfunction and infertility in a patient with 21-hydroxylase deficiency: effects of glucocorticoid treatment and surgery. *Exp Clin Endocrinol Diabetes* 2003;111(1):41-43.
101. Gao H-B, Tong M-H, Hu Y-Q, Guo Q-S, Ge R, Hardy MP. Glucocorticoid induces apoptosis in rat leydig cells. *Endocrinology*. 2002;143(1):130-138.

How to cite this article: Huang C, Ji X, Zhou W, et al. Coronavirus: A possible cause of reduced male fertility. *Andrology*. 2020;00:1-8. <https://doi.org/10.1111/andr.12907>