

Instructions.

- Vectors are indicated with arrows, as in

Let \vec{v} be a vector in \mathbb{R}^n .

- You are allowed a two-sided sheet of notes in your own handwriting.
- No calculators.
- There are 5 problems on 5 pages. Make sure your exam is complete.
- **Any cheating observed during, or noticed afterwards when comparing exams, will result in a 0 on this exam. Moreover, such an exam cannot be dropped. This means you lose 28% percent of your total raw grade. It will be difficult to pass the class if this occurs.**

Question	Points	Score
1	10	
2	12	
3	4	
4	15	
5	9	
Total:	50	

1. Check Yes or No for whether the subsets below are subspaces. You also need to EXPLAIN:

- If you check “Yes” explain why by doing one of the following: verifying the three properties, describing it as the span of a set of vectors, the nullspace of a matrix, the range of a linear transformation etc...
- If you check “No” give an explicit example of the property that is not satisfied.

[2 points] (a) The subset of \mathbb{R}^3 consisting of the vectors of the form $\begin{bmatrix} a \\ b \\ c \end{bmatrix}$ where $a - 3b = 5c$.
 Yes No Explanation:

[2 points] (b) The subset of \mathbb{R}^3 consisting of the vectors of the form $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ where $x + y + z = 1$.
 Yes No Explanation:

[2 points] (c) The set of all points on either the x -axis or the y -axis in \mathbb{R}^2 .
 Yes No Explanation:

[2 points] (d) The set of \vec{x} such that $T(\vec{x}) = \vec{0}$ where $T\left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}\right) = A^2 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$ and $A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 0 & 1 \\ 0 & -1 & -1 \end{bmatrix}$.
 Yes No Explanation:

[2 points] (e) The subset of \mathbb{R}^2 consisting of vectors of the form $\begin{bmatrix} a \\ b \end{bmatrix}$ where $a - b \leq 0$.
 Yes No Explanation:

[12 points] 2. Suppose that A is a 4×3 matrix, and that $\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 2 \\ 3 \end{bmatrix}$ are in the column space of A .

(a) Say if each of the following statements is true, false, or if it could be either true or false from the amount of information given.

1. The vector $A \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is in the column space of A .
 True False Not enough information
2. The vector $\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$ is in the column space of A .
 True False Not enough information
3. The nullspace of A is $\{\vec{0}\}$.
 True False Not enough information
4. The column space of A is $\{\vec{0}\}$.
 True False Not enough information
5. One of the columns of A is $\begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$.
 True False Not enough information

(b) Check off *all* values that the rank of A could be.

0 1 2 3 4

3. Let $\vec{u} = \begin{bmatrix} 1 \\ 3 \\ 2 \\ -1 \end{bmatrix}$ and $\vec{v} = \begin{bmatrix} 3 \\ 3 \\ 2 \\ 0 \end{bmatrix}$.

[2 points] (a) Write down a matrix you would row reduce to find two more vectors, \vec{x} and \vec{y} so that all four span \mathbb{R}^4 .

[2 points] (b) Do not row reduce! But, describe what you would look for in the echelon form of your matrix from the previous part to determine \vec{x} and \vec{y} .

4. Suppose you have the following matrices A and A^T that have echelon forms B and C .

$$A = \begin{pmatrix} 0 & 0 & 1 & 1 & -2 & 1 & 0 & 0 \\ -1 & 1 & -1 & 1 & 2 & 2 & -3 & 3 \\ -1 & 1 & 0 & 2 & 0 & 3 & -3 & 3 \\ 0 & 1 & -1 & 2 & 2 & 1 & -1 & 4 \\ -1 & 0 & 2 & 1 & -4 & 3 & -2 & -1 \end{pmatrix} \xrightarrow{\text{echelon}} B = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & -1 & 2 & 1 \\ 0 & 1 & 0 & 3 & 0 & 2 & -1 & 4 \\ 0 & 0 & 1 & 1 & -2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$A^T = \begin{pmatrix} 0 & -1 & -1 & 0 & -1 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & -1 & 0 & -1 & 2 \\ 1 & 1 & 2 & 2 & 1 \\ -2 & 2 & 0 & 2 & -4 \\ 1 & 2 & 3 & 1 & 3 \\ 0 & -3 & -3 & -1 & -2 \\ 0 & 3 & 3 & 4 & -1 \end{pmatrix} \xrightarrow{\text{echelon}} C = \begin{pmatrix} 1 & 0 & 1 & 0 & 2 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

In the following questions it is okay to use the notation \vec{a}_i for the i th column vector of A , or $\vec{a}_i(r)$ for the i th row vector. Similarly, for B and C . For A^T you can use \vec{a}_i^T and $\vec{a}_i^T(r)$.

[2 points]

(a) Find a basis for the row space of A that does NOT use any of the row vectors of A .

[2 points]

(b) Find a different basis for the row space of A that EXCLUSIVELY uses the row vectors of A .

[2 points]

(c) Find a basis for the column space of A that does NOT use any column vectors of A .

[2 points]

(d) Find a different basis for the column space of A that EXCLUSIVELY uses the column vectors of A .

Here are the matrices again:

$$A = \begin{pmatrix} 0 & 0 & 1 & 1 & -2 & 1 & 0 & 0 \\ -1 & 1 & -1 & 1 & 2 & 2 & -3 & 3 \\ -1 & 1 & 0 & 2 & 0 & 3 & -3 & 3 \\ 0 & 1 & -1 & 2 & 2 & 1 & -1 & 4 \\ -1 & 0 & 2 & 1 & -4 & 3 & -2 & -1 \end{pmatrix} \xrightarrow{\text{echelon}} B = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & -1 & 2 & 1 \\ 0 & 1 & 0 & 3 & 0 & 2 & -1 & 4 \\ 0 & 0 & 1 & 1 & -2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$A^T = \begin{pmatrix} 0 & -1 & -1 & 0 & -1 \\ 0 & 1 & 1 & 1 & 0 \\ 1 & -1 & 0 & -1 & 2 \\ 1 & 1 & 2 & 2 & 1 \\ -2 & 2 & 0 & 2 & -4 \\ 1 & 2 & 3 & 1 & 3 \\ 0 & -3 & -3 & -1 & -2 \\ 0 & 3 & 3 & 4 & -1 \end{pmatrix} \xrightarrow{\text{echelon}} C = \begin{pmatrix} 1 & 0 & 1 & 0 & 2 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

[2 points]

- (e) If the 5×9 augmented matrix $[A \mid \vec{e}_1]$ reduced to a consistent matrix then \vec{e}_1 is in the column space of A . Write down the SMALLEST dimension augmented matrix possible to determine if \vec{e}_1 is in the column space of A .

[2 points]

- (f) Without doing any further calculations, find the nullity of A^T .

[3 points]

- (g) Find a BASIS for the nullspace of A^T

5. Let S be the subspace of \mathbb{R}^3 given by vectors of the form $\begin{bmatrix} a \\ b \\ c \end{bmatrix}$ such that $a - 2b + 5c = 0$.

[2 points]

(a) Find a basis for S .

[2 points]

(b) Suppose $\{\vec{u}, \vec{v}, \vec{w}\}$ is a basis for \mathbb{R}^3 . Explain why a linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^n$ is completely determined by the values $T(\vec{u})$, $T(\vec{v})$, and $T(\vec{w})$.

[1 point]

(c) Find a linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$ such that the kernel of T is precisely S . By the previous part a satisfactory answer need only specify what it does to a basis for \mathbb{R}^3 .

Hint: You may use the fact that $\vec{w} = \langle 1, -2, 5 \rangle$ does not belong to S .

[2 points]

(d) Describe a linear transformation $F: \mathbb{R}^5 \rightarrow \mathbb{R}^3$ such that the range of F is S .

[2 points]

(e) What is the dimension of the kernel of such a linear transformation F as in part (d)?

Hint: it is not necessary to use your answer from part (d)