

# Introduction to Missing Data Analyses

2020 Summer Institute  
for the UC Adolescence Consortium

Craig Enders

University of California - Los Angeles

[cenders@psych.ucla.edu](mailto:cenders@psych.ucla.edu)

# Body Attitudes Data

Questionnaire data from a study of body attitudes in a sample of 500 middle school students

Variables include body mass index (BMI), five questionnaire items measuring negative body attitudes, and past history of being bullied (0 = never bullied, 1 = history of being bullied)

All questionnaire items measured on a 7-point scale

# Negative Body Attitudes Questionnaire

Strongly Disagree ←————→ Strongly Agree

---

1. My hips seem to broad to me.      1   2   3   4   5   6   7

2. I have a strong desire to be thinner.      1   2   3   4   5   6   7

3. I think I'm too thick.      1   2   3   4   5   6   7

4. Some parts of my body look swollen.      1   2   3   4   5   6   7

5. My belly looks as if I'm pregnant.      1   2   3   4   5   6   7

---

# bodyattitudes.dat

Variable	Name	Missing %	Scaling
Identifier variable	ID	0	Integer index
History of being bullied	BULLIED	10.4	0 = not bullied, 1 = bullied
Body mass index	BMI	8.0	Continuous
Attitude item 1 (hips too broad)	BATT1	13.4	7-point ordinal scale
Attitude item 2 (desire to be thinner)	BATT2	12.2	7-point ordinal scale
Attitude item 3 (too thick)	BATT3	0	7-point ordinal scale
Attitude item 4 (body looks swollen)	BATT4	12.4	7-point ordinal scale
Attitude item 5 (belly looks pregnant)	BATT5	0	7-point ordinal scale
Negative body attitudes composite	NEGBODYATT	29.6	Sum of five items


# Substantive Analysis

Negative body attitudes scale score regressed on body mass index and bullied indicator

$$NEGBODYATT_i = \beta_0 + \beta_1(BMI_i) + \beta_2(BULLIED_i) + \varepsilon_i$$

Substantive goal is to compare bullied and not bullied groups while controlling for BMI


# Regression Intercept



The intercept is the expected body attitudes score when bullied = 0

i.e., The mean for girls with no history of being bullied

# Regression Slope



The slope is the change in body attitudes score between the not bullied to bullied group

i.e., The mean difference for girls who have been bullied

# Missing Data Processes

The missing data process (mechanism) dictates the accuracy of a missing data handling method

Missing values could be haphazard and unrelated to the variables in the analysis model

Missingness is often systematic and could be related to observed data or the unseen values

# Observed And Missing Data

Hypothetically complete data

Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	3	5
7	1	6
2	1	6
5	9	3
3	2	2
1	6	7
9	4	9
2	5	6

=

Observed data

Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	NA	5
7	1	6
NA	1	6
5	9	3
3	NA	NA
1	6	7
9	4	9
2	NA	6

+


Missing data  
(latent, unseen)

Y	X <sub>1</sub>	X <sub>2</sub>
	3	
2		
	2	2
	5	

# Missing Completely At Random (MCAR)

Haphazard non-response,  
variables in the analysis do  
not predict missingness


e.g., Items missing due to a  
purely chance process



# Missing Completely At Random (MCAR)

Haphazard non-response,  
variables in the analysis do  
not predict missingness

e.g., Items missing due to a  
purely chance process



# Missing At Random (MAR)

Systematic non-response,  
observed scores in the  
analysis predict missingness


e.g., Observed variables  
such as BMI predict whether  
attitude items are missing

Y	X <sub>1</sub>	X <sub>2</sub>	Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3			
3		5	NA		
7	1	6			
	1	6	NA		
5	9	3			
3			NA	NA	
1	6	7			
9	4	9			
2		6	NA		

# Missing At Random (MAR)

Systematic non-response,  
observed scores in the  
analysis predict missingness


e.g., Observed variables  
such as BMI predict whether  
attitude items are missing



# Not Missing At Random (NMAR)

Systematic non-response,  
unseen (latent) scores in the  
analysis predict missingness


e.g., Attitude items are  
missing for participants with  
the most negative attitudes



# Not Missing At Random (NMAR)

Systematic non-response,  
unseen (latent) scores in the  
analysis predict missingness



e.g., Attitude items are  
missing for participants with  
the most negative attitudes





# Some Popular But Flawed Methods

Method	Properties
Exclude incomplete data records (completely or analysis-by-analysis)	Decreases power, requires unsystematic missingness, biased with systematic processes
Fill in missing values with the arithmetic mean	Atheoretical, severe bias under any process
Replace missing scores with predicted values from a regression equation	Distorts measures of variation and correlation
In longitudinal data, replace missing scores with the last observation from a preceding wave	Atheoretical, biased under any process



# Complete-Case Analysis (Deletion)



# Mean Imputation



# Regression Imputation




# Current Recommendations

Methods that assume systematic non-response due to observed scores (MAR) are often optimal

Easy to implement, widely available in software packages, and flexible for different applications

Maximum likelihood, Bayesian estimation, and multiple imputation assume an MAR process

# Relations Among Methods




## Practical Benefits

If missingness is related to observed scores ...


- a) Substantial missing data rates cause no bias
- b) Power is maximized because all available data contribute to the analysis
- c) Performance relative to other methods (e.g., deletion) improves as missing data rates increase

# Choosing A Missing Data Method




# Iterative Recipe For Missing Data Estimation

1) Estimate parameters, treating imputations as real data




2) Fill in missing values, treating parameters as true values


# Bivariate Scatterplot




# Distribution Of Attitudes Imputations




Imputation = Predicted Value + Noise




# Distribution Of BMI Imputations



Imputation = Predicted Value + Noise



# Imputed Data Set



# Blimp Bayes Regression Script

**DATA:** bodyattitudes.dat;

**VARIABLES:** id bullied bmi batt1 batt2 batt3 batt4 batt5 negbodyatt;

**NOMINAL:** bullied; ← **Categorical variable**

**MISSING:** 999; ← **Missing value code**

**MODEL:** negbodyatt ~ bmi bullied; ← **Regression model**

**CENTER:** bmi; ← **Center predictor**

**SEED:** 90291; ← **Random number seed (any integer)**

**BURN:** 2000; ← **MCMC iterations prior to analysis**

**ITERATIONS:** 10000; ← **MCMC iterations for analysis**

**OPTIONS:** psr; ← **MCMC diagnostic**

# Analysis Results

Parameter	Bayes		Maximum Likelihood		Multiple Imputation	
	Est.	S.D.	Est.	S.E.	Est.	S.E.
Intercept	19.780	0.209				
BMI	0.449	0.077				
BULLIED	4.487	0.638				

# Mplus FIML Regression Script

## DATA:

```
file = bodyattitudes.dat;
```

## VARIABLE:

```
names = id bullied bmi batt1 batt2 batt3 batt4 batt5 negbodyatt;
```

```
usevariables = bullied bmi negbodyatt; ← Select variables
```

```
missing = all(999); ← Missing value code
```

## DEFINE:

```
center bmi (grandmean); ← Center predictor
```

## MODEL:

```
bmi bullied; ← Predictor variables
```

```
negbodyatt on bmi bullied; ← Regression model
```

## OUTPUT:

```
stdyx; ← Standardized estimates
```

# Analysis Results

Parameter	Bayes		Maximum Likelihood		Multiple Imputation	
	Est.	S.D.	Est.	S.E.	Est.	S.E.
Intercept	19.780	0.209	19.747	0.198		
BMI	0.449	0.077	0.457	0.076		
BULLIED	4.487	0.638	4.652	0.646		

# Multiple Imputation Step 1: Create $M$ Complete Data Sets

Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	NA	5
7	1	6
NA	1	6
5	9	3
3	NA	NA
1	6	7
9	4	9
2	NA	6

Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	3.3	5
7	1	6
2.4	1	6
5	9	3
3	2.1	1.9
1	6	7
9	4	9
2	5.3	6

Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	4.7	5
7	1	6
1.3	1	6
5	9	3
3	6.5	3.5
1	6	7
9	4	9
2	4.2	6

Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	2.6	5
7	1	6
2.1	1	6
5	9	3
3	3.9	3.0
1	6	7
9	4	9
2	4.6	6

Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	3.3	5
7	1	6
2.4	1	6
5	9	3
3	2.1	1.9
1	6	7
9	4	9
2	5.3	6

Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	4.7	5
7	1	6
1.3	1	6
5	9	3
3	6.5	3.5
1	6	7
9	4	9
2	4.2	6

Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	2.6	5
7	1	6
2.1	1	6
5	9	3
3	3.9	3.0
1	6	7
9	4	9
2	4.6	6

1                      2                      ...                      M




# Multiple Imputation Step 2: Perform Analysis on Each Data Set

Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	3.3	5
7	1	6
2.4	1	6
5	9	3
3	2.1	1.9
1	6	7
9	4	9
2	5.3	6



Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	4.7	5
7	1	6
1.3	1	6
5	9	3
3	6.5	3.5
1	6	7
9	4	9
2	4.2	6

...


Y	X <sub>1</sub>	X <sub>2</sub>
4	4	3
3	2.6	5
7	1	6
2.1	1	6
5	9	3
3	3.9	3.0
1	6	7
9	4	9
2	4.6	6



## Multiple Imputation Step 3: Combine Estimates and Standard Errors



...



$$\hat{\theta} = (\hat{\theta}_1 + \hat{\theta}_2 + \dots + \hat{\theta}_M) / M$$

# Blimp Multiple Imputation Script

**DATA:** bodyattitudes.dat;

**VARIABLES:** id bullied bmi batt1 batt2 batt3 batt4 batt5 negbodyatt;

**NOMINAL:** bullied; ← **Categorical variable**

**MISSING:** 999; ← **Missing value code**

**FCS:** negbodyatt bmi bullied; ← **Variables to be imputed**

**SEED:** 90291; ← **Random number seed (any integer)**

**BURN:** 2000; ← **MCMC iterations prior to saving first data set**

**THIN:** 2000; ← **MCMC iterations between each additional data set**

**NIMPS:** 20; ← **Number of imputations**

**SAVE:** separate = imputation\*.dat; ← **Stack imputed data sets into a file**

**OPTIONS:** psr; ← **MCMC diagnostic**

# Mplus Analyze Imputations Script

## DATA:

file = imputationlist.dat; ← **Text file with 20 data set names**

type = imputation; ← **Imputation data**

## VARIABLE:

names = id bullied bmi batt1 batt2 batt3 batt4 batt5 negbodyatt;

usevariables = bullied bmi negbodyatt; ← **Select variables**

## DEFINE:

center bmi (grandmean); ← **Center predictor**

## MODEL:

negbodyatt on bmi bullied; ← **Regression model**

## OUTPUT:

stdyx; ← **Standardized estimates**

# Analysis Results

Methods are effectively equivalent!

Parameter	Bayes		Maximum Likelihood		Multiple Imputation	
	Est.	S.D.	Est.	S.E.	Est.	S.E.
Intercept	19.780	0.209	19.747	0.198	19.788	0.220
BMI	0.449	0.077	0.457	0.076	0.429	0.080
BULLIED	4.487	0.638	4.652	0.646	4.319	0.606

