




.....
Exoplanets
.....

Press Kit



Contents

Preface	3
Early discoveries	5
Techniques for detection	7
Direct detection	7
Imaging	7
Indirect detection	7
Radial velocity tracking	8
Astrometry	10
Pulsar timing	10
Transits	10
Gravitational microlensing	11
What can we learn from exoplanets?	13
What are exoplanets like?	14
Life outside the Solar System	16
Exoplanet research at ESO	17
ESO's current exoplanet instruments	18
Exoplanet research in the future at ESO	19



Cover: Artist's impression of the exoplanets HD 189733b | ESA, NASA, G. Tinetti (University College London, UK & ESA) and M. Kornmesser (ESO)

Left: Artist's impression of an exoplanets orbiting its star | ESA, NASA, M. Kornmesser (ESO) and STScI

Preface

Since planets were first discovered outside the Solar System in 1992 (orbiting a pulsar) and in 1995 (orbiting a “normal” star), the study of planets orbiting other stars, known as exoplanets, or extrasolar planets, has become one of the most dynamic research fields in astronomy. Our knowledge of exoplanets has grown immensely, from our understanding of their formation and evolution to the development of different methods to detect them.

This guide provides an overview of the history of exoplanets and of the current state of knowledge in this captivating field. It reveals the various methods that astronomers use to find new exoplanets and the information that can be inferred. The last section summarises the impressive findings of exoplanet research at ESO and the current and near-future technologies available in the quest for new worlds.



Early discoveries

“There are an infinite number of worlds, some like this world, others unlike it.”

Epicurus — letter to Herodotus
(~ 300 BC)

A planet is an object orbiting a star that is massive enough both to have achieved an almost spherical shape and to have cleared the rotating disc of dense gas, known as the protoplanetary disc, that surrounds a newly formed star. Planets differ in this from dwarf planets (such as Pluto), which do not have enough mass to clear the protoplanetary disc area.

The first detection of an exoplanet occurred in 1992 when the astrophysicists Aleksander Wolszczan and Dale Frail discovered three exoplanets. They were found in an unexpected environment, orbiting the pulsar PSR1257+12.

In 1995, the Geneva-based astronomers Michel Mayor and Didier Queloz detected the first exoplanet around a “normal” (main sequence) star, 51 Pegasi. The planet, named 51 Pegasi b, has around half the mass of Jupiter and whizzes around its parent star in just over four Earth days, lying almost eight times closer to it than Mercury is to the Sun.

Since 1995, this area of astronomy has become a very dynamic research field and astronomers have found over 450 exoplanets (as of May 2010), using a host of techniques.



Artist's impression of
the planetary system
around HD 69830 | ESO

Techniques for detection

Searching for exoplanets is like looking for the proverbial needle in a haystack. Planets emit little or no light of their own, while their host stars shine brightly. Seeing the light from a distant planet is like spotting a dim candle in front of a raging forest fire.

Nowadays six investigative tools are used to spot hidden exoplanets.

Direct detection

- Imaging

Indirect detection

- Radial velocity tracking
- Astrometry
- Pulsar timing
- Transits
- Gravitational microlensing

Direct detection

Imaging

The hardest way to detect an exoplanet is to try to image it directly. This is because of the extreme contrast between the light emitted by the parent star and by the companion planet. To expose the planet, the starlight must be dimmed or masked in some way so as to enable observers to see into the shadow. One method is to use infrared radiation, rather than visible light. The visible light output of a Jupiter-like planet is one billionth of that of its host star, while in the infrared the contrast is just a factor of a few thousandths. This is particularly true when the planet is still very young and thus contracting, thereby emitting heat. Another method is to physically block out the starlight, using a coronagraph that masks the bright central core of the star, leaving only the corona, the outer plasma region of the star's atmosphere, visible and so allowing any nearby planets to shine through.

Direct imaging is the only way to assess some important physical parameters, such as the amount of water on the surface and the properties of any possible biosphere.


The adaptive optics instrument, NACO, on ESO's Very Large Telescope (VLT) has obtained the first image of an exoplanet. The European Extremely Large Telescope (E-ELT) planned for 2018, will search for new planets using direct imaging, thanks to its very sharp vision.

Indirect detection


The majority of all exoplanets discovered so far have been detected using indirect methods — identifying their existence by their effect on their host star.


The presence of a planet affects its host star in several ways. The weak gravity of the planet pulls the star in a small circular orbit, introducing a minute wobble that can be detected using radial velocity tracking or astrometry (see pages 8–10). Alternatively, as the planet moves between the star and the observer, the measured luminosity of the star will change. These tiny variations are important for astronomers, as it makes the indirect observation of exoplanets possible.

Possibly the first image of an exoplanet (red spot), obtained with NACO at the VLT. The planet orbits a brown dwarf (blue spot in middle) | ESO



Beta Pictoris as seen in infrared light | ESO





The planetary system
around Gliese 581
(artist's impression) |
ESO


Radial velocity tracking

An astronomer can determine much about a distant star by recording its spectrum. As the star moves in the small orbit resulting from the pull of the exoplanet, it will move towards the Earth and then away as it completes an orbit. The velocity of the star along the line of sight of an observer on Earth is its radial velocity. Changes in the radial velocity of the star cause the lines in the star's spectrum to shift towards redder wavelengths when the star is moving away from us and towards bluer wavelengths when the planet is approaching us (see image). This is the Doppler effect, and it is noticeable with sound waves in everyday life,


for example in the change of pitch of an ambulance siren as it drives past on the street.


The periodic changes in the star's radial velocity depend on the planet's mass and the inclination of its orbit to our line of sight. These tiny changes or "wobbles" can be measured by a distant observer. Astronomers use high precision spectrographs to study Doppler-shifted spectra, looking for small regular variations in the radial velocity of a star. As the inclination of the planetary orbit is unknown, the measurement of this regular variation gives a minimum value for the mass of the planet.

The radial velocity method has proven to be the most successful in finding new planets. At present, the most successful low-mass exoplanets hunter is HARPS (High Accuracy Radial Velocity for Planetary Searcher), which is mounted on ESO's 3.6-metre telescope at La Silla, Chile.




The radial velocity method | ESO







1. The measured drop in brightness of the star when the planet passes in front of it | ESO



2. Brightness variations of two stars with transiting exoplanets | ESO



3. Light curve of OGLE-2005-BLG-390 | ESO



4. Gravitational lensing caused by the presence of a star and an exoplanet | ESO

Astrometry

The astrometry method is similar to radial velocity tracking and is used to detect exoplanets by measuring the small regular perturbation in the position of a star due to its unseen companion. The star moves in a tiny circular orbit on the sky with a radius that depends on the mass of the planet and its distance from the star, but not on the inclination. No planets have been discovered so far using this method.

Pulsar timing

The presence of a planet orbiting a star affects the timing of the regular signals emitted by the star itself. This phenomenon can be used to detect planets around a pulsar. Pulsars emit radio waves regularly as they rotate, creating a periodically pulsed beam, like a lighthouse. If an orbiting planet perturbs the motion of the star, then the timing of the beam is also affected, and this is how the first exoplanets were detected.

Transits

When a planet passes between the Earth and its host star, this is known as a transit. The planet blocks some of the starlight during the transit and creates a periodic dip in the brightness of the star. This effect can be measured using photometry, which measures the amount of light coming from celestial objects.

We can learn much about the composition of a planet's atmosphere from planetary transits. As a planet passes in front of its star, light from the star will pass through the planet's atmosphere, where some of it is selectively absorbed. By comparing the "before" and "after"



The Crab Nebula | ESO

spectral data of the starlight, the composition of the planet's atmosphere can be deduced.

The Optical Gravitational Lensing Experiment (OGLE) located at Las Campanas, Chile, was used to find the first planet through transit photometry (called OGLE-TR-56). Nowadays, satellites in space such as COROT and Kepler have found numerous transiting planets.


Radial velocity measurements, combined with transit photometry, make it possible to determine not only the mass of a planet, but also its radius and density.

Gravitational microlensing

The gravitational pull of a large object will bend the light from distant objects and amplify it, acting like a magnifying lens. When light from the background object travels towards Earth, its path is bent or warped as it bypasses any large foreground object that is aligned with the background light source. As the microlensing effect works on radiation from the background source, this technique can be used to study intervening objects that emit little or no light, such as black holes, or planets around distant stars. Suppose that the aligned foreground mass to be studied

is a star that is hosting a planet, then the amplified light curve from the background source will contain an additional side peak. The size and shape of the secondary peak will depend on the mass and distance of the planet from the host star (see the image).


The exoplanet OGLE 2003-BLG-235/MOA 2003-BLG-53 was the first planet discovered using this technique, in 2003. The disadvantage of the microlensing technique is that the effect happens only once, as it relies on a unique chance alignment of the foreground and background stars, and so measurements must be checked using other methods.



Planet-forming disc
(artist's impression) |
ESO

The central region of the
Orion Nebula | ESO,
M. McCaughrean et al.

Possible orbital
migration of the plane-
tary system around
HD 89830. Planets may
have formed far away
from the star and spi-
ralled inwards over time. |
ESO



What can we learn from exoplanets?

Exoplanets are fascinating because they may solve mysteries about our own Solar System. There is a wealth of data available to study different types of galaxies and stars, which has enabled astronomers to develop models and theories on star and galaxy formation and to place our own galaxy and star amongst them. The Solar System is 4.6 billion years old, but there is no way to measure directly how it formed and it was, until recently, the only planetary system that we knew of, so there was nothing to compare it with. We had no idea if it was one of many, a typical example of a planetary system or a unique one-off. Studying the formation of other young planetary systems may give us answers.

Protoplanetary discs are regions of dust and gas orbiting very young stars, where planets are formed. Current theories of planetary formation suggest that dust particles start to collapse under gravity and stick together, forming bigger and bigger grains. If young protoplanetary discs survive the threat of stellar radiation and impacts by comets and meteorites, then matter continues to clump together and eventually planetoids may form. Planetoids are celestial objects bigger than meteorites and comets, but smaller than planets. After a few million years, most of the circumstellar dust will have been swept away as planetoids accumulate mass and grow into planets.


Many of the planets found so far are large, gaseous and very close to their star, unlike the situation in our own Solar System. The concept of orbital migration has been revived to explain the close proximity of some giant planets to their star: these planets may have formed undisturbed relatively far from the star and then slowly spiralled inwards over time.



What are exoplanets like?


Due to the limitations in current detection methods, the majority of planets discovered so far have been rather large — Jupiter-sized or much larger. Although it is difficult to detect smaller planets, a planet with less than twice the mass of the Earth has been discovered.


There are small icy exoplanets as well as gigantic hot planets. One of the interesting questions to answer is how the distribution of exoplanet type is linked to the type of parent star. It is likely that there are also exoplanets with rings and satellites, but these are difficult features to detect.



This diagram compares our Solar System with some of the various planetary systems imaged so far (the Solar System is at the bottom of the image, showing the Sun along with the four outer planets, Jupiter, Saturn, Uranus and Neptune)

Artist's concept of the exoplanet orbiting Fomalhaut | ESO (L. Calçada), ESA, NASA





Artist's impression of an exoplanetary system | NASA, ESA and G. Bacon

Life outside the Solar System

The current focus of research into exoplanets is to develop the theories and understanding of planetary formation, and to understand how the Solar System developed and what its future might be. However, what makes exoplanets truly fascinating for most is the possibility of finding another world that harbours life.

Exobiology is concerned with the study of life outside of the Earth. The concept of "life" is subject to debate, but there is agreement in defining the features that could permit the development of carbon-based life:

- A planet should have a mass of between 1 and 10 terrestrial masses, be big enough to hold its atmosphere, but not so massive that it keeps too much hydrogen.
- A planet must be in the habitable zone, sometimes called the Goldilocks Zone, which is defined as the band around a star where water can be liquid. This means that a planet can neither be too close nor too far from its star, as water would be either gaseous or icy respectively.

Exobiology is not a focus of current exoplanet research projects, but is one for the future. Future spectroscopic missions — ESA's Darwin and NASA's Terrestrial Planet Finder missions — are planned for launch over the next decade and will search for oxygen, carbon dioxide and chlorophyll.

Exoplanet research at ESO

A list of ESO's most recent achievements is given below.

- 2010: VLT detects first superstorm on an exoplanet. (eso1026)
- 2010: For the first time, astronomers have been able to directly follow the motion of an exoplanet as it moves to the other side of its host star. (eso1024)
- 2010: Six exoplanets were found orbiting in the opposite direction to the rotation of their host star — challenging theories of planet formation. (eso1016)
- 2010: With HARPS, astronomers have discovered the first “normal” exoplanet that can be studied in great detail. (eso1011)
- 2010: VLT captures first direct spectrum of an exoplanet. (eso1002)
- 2009: Astronomers discover first super-Earth with an atmosphere. (eso0950)
- 2009: Sun-like stars that host planets appear to have destroyed their lithium much more efficiently than “planet-free” stars. (eso0942)
- 2009: HARPS discovers 32 new exoplanets, mostly low-mass ones. (eso0939)
- 2009: HARPS finds first solid evidence for a rocky exoplanet. (eso0933)
- 2009: Lightest exoplanet found using the most successful low-mass exoplanet hunter in the world, the HARPS spectrograph. (eso0915)
- 2008: First planet discovered around a fast-rotating hot star, discovered by three undergraduate students and confirmed by ESO's VLT. (eso0845)
- 2008: First direct image of a planet that is as close to its host star as Saturn is to the Sun. (eso0842)
- 2008: Unsurpassed details revealed on the motion and makeup of planet-forming discs around Sun-like stars. (eso0827)
- 2008: A trio of super-Earths are observed using ESO's HARPS instrument. Data suggests one in three Sun-like stars have such planets. (eso0819)
- 2007: Discovery that exoplanets may pollute the atmospheres of their parent stars with planetary debris. (eso0729)
- 2007: ESO develops a new imaging spectrograph so as to be able to image faint objects obscured by their bright parent stars directly. This paves the way for many thrilling new discoveries. (eso0728)
- 2007: Discovery of the most Earth-like planet: located 20 light-years away, it may have water on its surface. (eso0722)
- 2006: Observations show that some objects that are several times the mass of Jupiter have a disc surrounding them and may form in a similar way to stars. It thus becomes much more difficult to define precisely what a planet is. (eso0629)
- 2006: Detection of three Neptune-like planets, each of a mass between ten and twenty times that of Earth, around a star that also possesses an asteroid belt. Of all known systems, this is the most similar yet to our own Solar System. (eso0618)
- 2006: Discovery of the first terrestrial-sized exoplanet, five times the size of the Earth. (eso0603)
- 2005: Discovery of a planet with a mass comparable to Neptune around a low-mass star, the most common type of star in our galaxy. (eso0539)
- 2004: Ingredients for the formation of rocky planets discovered in the innermost regions of protoplanetary discs around three young stars. This suggests that the formation of Earth-like planets may not be unusual. (eso0435)
- 2004: First direct image taken of an exoplanet, paving the way for more direct studies. (eso00428)
- 2004: Discovery of the first possible rocky exoplanet, an object with 14 times the mass of the Earth. (eso0427)
- 2004: Confirmation of the existence of a new class of giant planet. These planets are extremely close to their host stars, orbiting them in less than two Earth days, and are therefore very hot and “bloated”. (eso0415)
- 2002: The discovery of a dusty, opaque disc surrounding a young Sun-like star, in which planets are forming or will soon form. This disc is similar to the one in which astronomers think the Solar System formed. (eso0214)

The VLT platform at
Paranal | ESO




ESO's current exoplanet instruments

The groundbreaking discoveries of recent years were possible thanks to the ESO instruments searching for following:

- HARPS (High Accuracy Radial Velocity for Planetary Searcher) on the ESO 3.6-metre telescope at La Silla for radial velocity high resolution spectroscopy. It can measure velocities with a precision greater than 1 m/s (or 3.6 km/h).
- NACO on the VLT at Paranal — an active optics and near-infrared imager and spectrograph that allows the sharp imaging of objects smaller and fainter than stars, such as exoplanets.
- UVES on the VLT — for radial velocity high resolution spectroscopy in the UV and visible.
- EMMI on the NTT at La Silla — spectrograph operating at visible wavelengths.
- FLAMES (Fibre Large Array Multi Element Spectrograph), on ESO's VLT at Paranal — for multi-fibre resolution spectroscopy.
- Swiss 1.2-metre Leonhard Euler Telescope at La Silla — high resolution spectroscopy.
- Danish 1.54-metre Telescope at La Silla — long-term monitoring.
- AMBER studies of circumstellar environments and protoplanetary discs, important for planet formation studies.
- VISIR studies of circumstellar environments and protoplanetary discs, important for planet formation studies.
- ISAAC (Infrared Spectrometer And Array Camera), attached to the VLT
- MIDI, the MID-infrared Interferometric instrument of the VLT Interferometer.

The Atacama Large Millimeter/submillimeter Array | ALMA (ESO/NAOJ/NRAO)/L. Calçada (ESO)





The E-ELT | ESO


Exoplanet research in the future at ESO

The PRIMA instrument of the ESO Very Large Telescope Interferometer (VLTI), which recently saw “first light” at its new home atop Cerro Paranal in Chile, will boost the capabilities of the VLTI to see sources much fainter than any previous interferometers, and enable astrometric precision unmatched by any other existing astronomical facility. PRIMA will therefore be a unique tool for the detection of exoplanets.

The second generation instrument SPHERE for ESO’s Very Large Telescope is dedicated to the discovery and study of new giant exoplanets orbiting nearby stars by direct imaging, in particular of planets more massive than Jupiter at various stages of their evolution, in the key separation regime 1 to 100 AU (AU is the mean Earth–Sun distance). SPHERE should have first light around 2011.

Moreover, two future ground-based telescopes will be used to search for exoplanets:

- E-ELT (European Extremely Large Telescope) — expected to be able to image exoplanets directly, revealing their composition, and to detect, via the radial velocity method, Earth-mass planets.
- ALMA (Atacama Large Millimeter/submillimeter Array) — for accurate astrometry measurements, possibly even for direct detection. Detailed mapping of protoplanetary discs, which is important for understanding planet formation.



Exoplanets

Press Kit

