

CHEMISTRY MODEL QUESTION PAPER (3rd SET SOLUTION) SSC-I

SECTION A

Q1: MCQs

1	2	3	4	5	6	7	8	9	10	11	12
D	C	A	D	C	A	D	A	D	C	В	A

SECTION – B (Marks 33)

Q.2 Attempt any ELEVEN parts from the following. All parts carry equal marks.

$$(11 \times 3 = 33)$$

i. Differentiate between analytical and physical chemistry (at least two).

Answer:

Analytical Chemistry

It is the branch of chemistry which deals with qualitative and quantitative analysis of substances. For example, study of soil components.

Physical Chemistry:

Physical Chemistry is the branch of chemistry that deals with the laws and theories to understand the structure and physical and chemical changes of matter. Such as laws of gases, properties of solids etc.

ii. Explain the method of preparation of 0.5M NaOH in 100cm³ solution from 1M NaOH.

Answer: Given Data

Molarity of stock solution of NaOH= $M_1 = 1M$

Molarity of dilute solution of NaOH to be prepared = $M_2 = 0.5 \text{ M}$

Volume of the dilute solution of NaOH = 100cm^3

Volume of the stock solution = V1 = ?

Using dilution formula, a dilute solution can be prepared from a stock solution.

Therefore


$$M_1 \ x \ V_1 \ = \ M_2 \ x \ V_2$$

$$V_{1} = M_{2} \times V_{2} = 0.5M \times 100 \text{cm}^{3} = 50 \text{ cm}^{3}$$
 $M_{1} = 1M$

Thus, it is evident that by taking 50 cm³ of the stock solution and transferring it to a volumetric flask of 100 cm³ and making it up to the mark give rise to 0.5M dilute solution of NaOH.

iii. Draw the structure of isotopes of chlorine.

Answer: Isotopes of Chlorine



iv. Briefly explain octet and duplet rule with example.

Answer:

Duplet Rule


The tendency of atoms to acquire two electron configurations in their valence shell during chemical bonding is called duplet rule.



Each hydrogen share one electron so in a molecule it becomes two electrons.

Octet Rule:

The tendency of atoms to acquire eight electron configurations in their valence shell during chemical bonding is called octet rule. Most of the elements have incomplete octet, they form bond to complete their octet such as Nitrogen needs three electrons to complete its octets. So each N atom shares three unpaired electrons with another one to complete its octet.



v. Identify the characteristic of ionic compounds.

Answer: Properties of Ionic Compounds

- a) Ionic bond is a very strong bond, therefore, ionic compounds have high melting point.
- b) Ionic compounds form crystals.
- c) Ionic compounds do not conduct electricity in a solid-state but they do conduct electricity in the molten state.

vi. Demonstrate diffusion and effusion of the gasses with the help of examples.

Answer:

Diffusion:

Diffusion is defined as spontaneous mixing up of molecules by random motion to form a homogeneous mixture. Diffusion occurs due to difference in concentration.

Examples: We can smell perfume because it diffuses into the air.

Effusion:

The escape of molecules of gases through a tiny hole is called effusion. For example when a tyre gets punctured, air effuses out.

vii. Differentiate between saturated and unsaturated solutions (at least two).

Answer:

S.No	Unsaturated Solution	Saturated Solution
01	Unsaturated solutions are solutions in	Saturated solutions are the solutions in
	which the amount of dissolved solute is	which the amount of dissolved solute at a
	less than the saturation point of the solvent	specific temperature is equal to the
	at a specific temperature.	saturation point of the solvent.
02	More solutes can be dissolved at the same	More solutes cannot be dissolved at the
	temperature.	same temperature.

viii. Describe the formation of solution by mixing solid into gases with example.

Answer: Solid particles released into the air get mixed with air forming a solid-gas solution. For example, carbon particles from different sources released into the air such as smoking.

ix. State the common rules for assigning the oxidation number.

Answer:

Common Rules For Assigning Oxidation Number

- a. The oxidation number of elements in a free or uncombined state is always zero. For example, the oxidation state of all free state elements like Cl_2 , H_2 , Zn, Na, and Mg, etc are zero.
- b. The oxidation number of a single ion is the same as the charge on ions. For example, the oxidation states of Na^+, Ca^{+2}, Cl^- , Al^{3+} , and S^{2-} are (+1), (+2), (-1), (+3), and (-2) respectively.
- c. The oxidation number of hydrogen in all of its compounds is (+1), but if hydrogen is linked with metals, then the oxidation state of hydrogen is seen as (-1). For example, in HCl, the oxidation number of hydrogen is (+1).
- d. The oxidation number of each of the atoms in a molecule is counted individually and their algebraic sum is zero. For example, In KClO₃. Using formula:

KClO₃

$$+1 + x - 2 \times 3 = 0$$

$$X = 5$$

Hence, oxidation state of Cl is +5. And of KClO₃ is zero.

x. List three uses of electrolytic cells.

Answer:

Uses of electrolytic cell

- a. Electrolysis is used in the extraction of metals from their ores.
- b. It is used for refining certain metals such as copper and zinc.
- c. Electrolysis is used for the manufacture of chlorine gas.
- d. Electrolysis is used for electroplating many things we use every day.

xi. Write down the oxidation and reduction reaction in voltaic cell at Anode and Cathode.

Answer:

Reactions in Voltaic Cell Reaction at the Anode:

Zn/ZnSO₄ half cell, oxidation reaction occurs at the anode.

$$Zn^{\circ}$$
 $Zn^{2+} + 2e^{-}$ (oxidation)

Reaction at the Cathode

Cu/CuSO4 half-cell, reduction reaction occurs at the cathode.

$$Cu^{2+} + 2e^{-}$$
 Cu° (reduction)

xii. Show how cations and anions are related to term metals and nonmetals.

Answer:

Metals have least number of valence electrons. Therefore, they always tend to lose electron forming cations. For example, Na has only one electron in its valence shell, so it will lose only one electron.

$$Na \longrightarrow Na^+ + 1e^-$$

However, non-metals require least number of electrons to fill their valence shell, therefore, when they acquire electron (s), they form anions.

xiii. Briefly describe why alkali metals are not found in free state in nature.

Answer: Alkali metals are highly reactive because of single valence electron with least ionization energy value. For example, if sodium metal is exposed to air, it reacts violently with the oxygen present in the air forming sodium oxide. That is the reason that Na is preserved in Kerosene oil to avoid contact with the air. Thus it is clear that all alkali metals form bonds with other elements to acquire stability and are not found in free state naturally.

xiv. Tabulate soft and hard metals with suitable examples.

Answer:

Soft Metals	Hard Metals
Soft metals are those metals which can be cut	Hard metals are those which cannot be
and easy to dent, work, or cut without shattering	easily cut. Since hardness is the
and affecting its malleability.	resistance to deformation. They exhibit
Sodium metal is soft enough to be cut with a	signs of high density and a tolerance of
knife. Potassium is also a soft metal. The	high temperatures. Therefore, these
cohesive forces between atoms of these metals	metals have stable physical formation.
are very weak. Thus the metals have a soft	Metallic bond in such metals is
texture.	stronger as compare to the soft metals.
Other examples of soft metal include gold,	Examples of hard metals are Tungsten,
silver, aluminum, lead etc.	Iridium, Chromium, Osmium.

xv. List the commercial value of silver, gold and platinum.

Answer:

Uses of Gold

It is used as an ornamental metal, gold coins etc.

Uses of Silver

Alloys of silver with copper are widely used in making coins, silver-ware and ornaments.

Uses of Platinum

An alloy of platinum, palladium and rhodium is used as catalyst in automobiles as catalytic convertor.

SECTION - C

Q.3 a. Differentiate between oxidation and reduction in term of oxygen and hydrogen with equations.

Answer:

Oxidation	Reduction
Oxidation can be defined as addition of	Reduction can be defined as addition of
Oxygen.	Hydrogen.
2Cu + O ₂ → 2CuO	$Cl_2 + H_2 \rightarrow 2HCl$
Copper is oxidized in above reaction.	Chlorine is reduced in above reaction.
Oxidation can also be defined as removal	Reduction can also be defined as removal of
of Hydrogen.	Oxygen.
$CH_4 + H_2 \rightarrow CO_2 + H_2O$	$CuO + H_2 \rightarrow Cu + H_2O$
Carbon is oxidized in above reaction.	Copper is reduced in above reaction.

Q3 (b) Identify the relationship between electronic configuration and the position of an element in the periodic table. $_{19}K^{39},_{17}Cl^{35}$ and $_{16}S^{32}$.

Answer:

$_{19}K^{39}$

Electronic Configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 4s^1$ Total No. of Shells = 4 It belongs to period 4. Total No. of Electrons in valance shell = 1 It belongs to group I-A.

17Cl³⁵

Electronic Configuration: $1s^2 2s^2 2p^6 3s^2 3p^5$ Total No. of Shells = 3 It belongs to period 3. Total No. of Electrons in valance shell = 7 It belongs to group VII-A.

$_{16}S^{32}$

Electronic Configuration: $1s^2 2s^2 2p^6 3s^2 3p^4$ Total No. of Shells = 3 It belongs to period 3. Total No. of Electrons in valance shell = 6 It belongs to group VI-A.

Q4 (a) Define empirical and molecular formula. Show the formation of empirical formula from molecular formula of the given compounds: $C_6H_{12}O_6$, $C_8H_{16}O_2$ and $C_{12}H_{22}O_{11}$.

Answer:

Empirical Formula:

Formula showing simple whole number ratio of atoms in the compound is called empirical formula. **Molecular Formula:**

Formula showing actual number of atoms in any compound is called molecular formula.

i) Molecular Formula: $C_6 H_{12} O_6$

C: H:O 6:12:6 1:2:1

Empirical Formula: C H₂ O

ii) Molecular Formula: $C_8 H_{16} O_2$

C: H:O 8:16:2 4:8:1

Empirical Formula: C₄ H₈ O

iii) Molecular Formula: $C_{12} H_{22} O_{11}$

C: H:O 12:22:11

(Cannot be simplified any further)

Empirical Formula: C₁₂ H₂₂ O₁₁

Q4 (b) Compare the physical state of matter with regards to intermolecular forces between them.

Answer:

SOLID	LIQUID	GAS	
Particles have strong	Particles have intermediate	Particles have weak	
intermolecular forces.	intermolecular forces.	intermolecular forces.	
Due to strong	Due to intermediate	Due to weak intermolecular	
intermolecular forces,	intermolecular forces,	forces, particles are far away.	
particles are very	particles are little far.		
close.			
Due to strong	Due to intermediate	Due to weak intermolecular	
intermolecular forces,	intermolecular forces,	forces, particles move	
particles have fixed	particles do not have fixed	randomly in all directions.	
position.	positions.		

Q5 (a) Use the rule that "like dissolves like" Describe dissolution of KCl in water with the help of diagram.


Answer:

When KCl is put in water, it immediately splits into K^+ and Cl^- .

$$KCl \rightleftharpoons K^+ + Cl^-$$

Water, H_2O , is a polar molecule, and has partial positive $^+\delta$ and negative charges $^-\delta$. The hydrogens on water are partially positive, and the oxygen on water is partially negative.

Since opposite charges attract each other. So the oxygen of the water molecules are attracted to the K^+ , and the hydrogens are attracted to the Cl^- . The water actually surrounds the K^+ and Cl^- so that the opposite charges are allowed to be close together to form hydrate and process is called hydration (solvation). That is why KCl is dissolved in water.



Q5 (b) How will you discuss the reactivity of halogens by using following reactions:

i)
$$KI + Br_2$$
 \longrightarrow $2KBr + I_2$
ii) $KBr + Cl_2$ \longrightarrow $2KCl + Br_2$

ii)
$$KBr + Cl_2 \longrightarrow 2KCl + Br_2$$

Answer:

$$F_2 > Cl_2 > Br_2 > I_2$$

Oxidizing power of F_2 is highest and that of I_2 is lowest.

$$KI + Br_2 \rightarrow 2KBr + I_2$$

 I_2 cannot oxidize any of halide ion whereas Br_2 can oxidize Γ^1 ion.

$$KBr + Cl_2 \rightarrow 2KCl + Br_2$$

 Cl_2 is a better oxidizing agent than Br_2 . It can oxidize both Br^{-1} and Γ^{-1} .