

Comprehensive Geriatric Assessment

A Summary



*Adapted from Specialist Geriatric Team Guidance on Comprehensive
Geriatric Assessment
National Clinical Programme for Older People
(2016)*

Overview

Comprehensive Geriatric Assessment (CGA) is fundamental to the assessment, planning and intervention required to meet the health and social care needs of the older person that is frail or at risk of frailty. Rather than the traditional way of working separately, CGA results in doctors, nurses, physiotherapists, occupational therapists, social workers and other members of the team working closely together to ensure an integrated assessment and response to the older person's individual needs. CGA has the potential to improve the care they receive in hospital, reduce unnecessary hospital admissions, lengths of stay and readmissions.

What is a Comprehensive Geriatric Assessment?

Comprehensive geriatric assessment (CGA) is an organised approach to assessment designed to determine an older person's medical conditions, mental health, functional capacity and social circumstances. Its purpose is to develop and implement a coordinated and integrated plan for treatment, rehabilitation, support and long term follow up. CGA is based on the premise that a full evaluation of a frail older person by a team of healthcare professionals may identify a variety of treatable health problems, resulting in a co-ordinated plan and delivery of care leading to better health outcomes.

Substantial evidence shows that in hospital, those who receive inpatient CGA on specialist geriatric wards are more likely to return home, are less likely to have cognitive or functional decline and have lower mortality rates than those who are admitted to general wards (*Ellis et al., 2011*).

Benefits of CGA

The benefits of CGA, in comparison to less structured multidisciplinary assessment where each discipline approaches the patient's assessment and plan of care in isolation, include:

- improves diagnostic accuracy
- optimises medical and rehabilitation treatment
- enhances health and functional outcomes
- informs the development of individualised care plans
- assists in avoiding the potential complications of hospitalisation
- facilitates effective discharge planning.

Who should have CGA?

The National Clinical Programme for Older People recommends that all older adults identified as being frail or at risk of frailty should have a timely comprehensive geriatric assessment performed and documented in their permanent health record (HSE 2012).


Note on Frailty

Older people living with frailty or at risk of frailty are admitted to hospital more frequently, have longer length of stay and occupy more bed days in acute hospitals in comparison to other patient groups.

Frailty is increasingly recognised as a distinctive state of health related to the aging process in which multiple body systems gradually lose their inbuilt reserves. It is not an inevitable part of ageing; it is a long term condition in the same sense as diabetes or Alzheimer's disease (BGS 2014). Clinically, older people who are frail have poor functional reserve, so that even a relatively minor illness or a change in medication can lead to a sudden catastrophic functional decline – causing the person to fall, become immobile or rapidly confused, or to present non-specifically with failure to thrive (Clegg et al 2013).

Frailty is a graded abnormal health state which ranges from mildly frail (those who need supported self-management), through those who are moderately frail and would benefit from structured interventions from healthcare professionals, to those who have advanced frailty where anticipatory care planning and end-of-life care may be appropriate interventions. Frail people should not be perceived as a problem to the system but, rather, clinicians should support people living with frailty to maintain their own health for as long as possible (NHS 2014).

Pattern of Frailty



Identifying Frailty

There are two possible approaches to identifying frailty:

Presence of frailty syndromes:

Falls (e.g. collapse, legs gave way, 'found lying on floor')

Immobility/ decreased mobility (e.g. sudden change in mobility, 'gone off legs', 'stuck in toilet')

Delirium (e.g. acute confusion, 'muddled', sudden worsening of confusion in someone with previous dementia or known memory loss)

Incontinence (e.g. change in continence – new onset or worsening of urine or faecal incontinence)

Susceptibility to side effects of medication (e.g. confusion with codeine, hypotension with antidepressants) (British Geriatric Society 2014).

Validated ***frailty assessment tools:***

Rockwood Clinical Frailty Scale.

PRISMA 7 Questionnaire

Timed up and go Test

The Groningen Frailty Indicator Questionnaire

Edmonton Frail Scale

When is CGA indicated?

When an older person is identified as being at risk of frailty, whether in an acute hospital, day hospital, community or residential care, they should be considered for a CGA. The content of the assessment may vary depending on different settings of care (e.g., home, clinic, hospital, nursing home). A key element of CGA is that the environment in which it is delivered is gerontologically attuned (McCabe & Kennelly 2015).

If indicated, CGA should be initiated as soon as possible after admission to hospital by a skilled, senior member of the multidisciplinary team, and used to identify reversible medical problems, target rehabilitation goals and plan all the components of discharge and post-discharge support needs (Oliver *et al* 2014).

Who should carry out CGA?

Members of the CGA multidisciplinary core team should include experienced individuals drawn from medical, nursing and health and social care professions. This multidisciplinary team is responsible for the coordinated assessment, discussion and recommendation or implementation of treatment plans.

The CGA **multi-disciplinary team** may include:

- Medical – e.g. geriatrician, psychiatry of old age, palliative care specialist, GP
- Nursing
- Medical social worker
- Physiotherapy
- Occupational therapy
- Speech and language therapy
- Dietetics
- Pharmacists
- Podiatry

Many components of CGA can be jointly undertaken by a number of disciplines on the team as some of the assessments and interventions overlap.

Expertise from other disciplines may be required for specific aspects of assessment.

Other team members may include podiatry, psychology, psychiatry of old age or other areas of speciality.

In most teams, a senior geriatrician will have the responsibility for team leadership. Experienced team members can vary in their roles including physical and functional assessments, providing advice and coordination of teams and services. The geriatrician and medical team often serve an integrative function, putting together the findings from other disciplines, and interpreting results of investigations in order to find a unifying diagnosis and hence guide a comprehensive, interdisciplinary management and treatment plan.

Elements of CGA

No one size fits all – CGA can be developed locally depending on the members and seniority of the multidisciplinary team. The assessment should, however, consist of the elements depicted here:

The four main dimensions of CGA are listed overleaf.



Four main dimensions of assessment

The four main dimensions covered in a CGA should include physical, functional, psychological and social assessment as follows:

Physical assessment <ul style="list-style-type: none">• Presenting complaint• Past medical history• Medication reconciliation and review• Nutritional status• Alcohol• Immunisation status• Advanced directives	Functional assessment <ul style="list-style-type: none">• Activities of daily living• Balance• Mobility
Psychological assessment <ul style="list-style-type: none">• Cognition and mood	Social assessment <ul style="list-style-type: none">• Living arrangements• Social support• Carer stress• Financial circumstances• Living environment

References

British Geriatric Society. (2014). *Fit for Frailty: Consensus best practice guidance for the care of older people living with frailty in community and outpatient settings*. British Geriatric Society: London.

Clegg, Dr. A., (2013) *Frailty in Elderly People*. The Lancet, Volume 381, Issue 9882, 8-14 page 1328

Ellis, G., Whitehead, M A., O'Neill, D., Langhorne, P., Robinson, D. (2011) *Comprehensive geriatric assessment for older adults admitted to hospital: meta-analysis of randomised controlled trials*. BMJ, 343: d6553.

HSE. (2012). *National Clinical Programme Older People. Specialist Geriatric Services Model of Care Part 1: Acute Service Provision*.

HSE. (2015). *National Clinical Programme Older People. (2015) Specialist Geriatric Team Guidance on Comprehensive Geriatric Assessment*.

NHS. (2014) *Safe, compassionate care for frail older people using an integrated care pathway: Practical guidance for commissioners, providers and nursing, medical and allied health professional leaders*.

McCabe, J., Kennelly, S. (2015). Acute care of older patients in the emergency department: strategies to improve patient outcomes. *Open Access Emergency Medicine* ,7, 45–54

Oliver, D., Foot, C., Humphries, R. (2014) *Making our health and care systems fit for an ageing population*. Kings Fund