

health is an idiot!" and propose him out of the post he held (in the Department of Agriculture) little wonder that it was not bring a degree of prestige to. Because of restraints to induction, the only role it ever attained of policing, and, as we policeman's lot is not a happy

reminded of that tragic character only on his deathbed, realized he never lived at all. The FDA is a sadder case; it never lived and died. We have written this as a warning because we believe that good can come from a public realization that this is, to all intents and purposes,

the only proper for the Congress to remove the FDA with full military might a gallant and futile battle on possible odds. More than that, it is useful for the Congress to take note of the fact that the office no longer has that it may order the formation of an agency empowered to do what it is charged to do, tried to do, or permitted to do. Now is the time to sign a Federal Drug Commission bill from a political superstructure that warms and emasculates it.

Walter Modell, M.D.

On December 12 that Commissioner of Food and Drugs. His job must have been. We

The Editor

Commentary

The evaluation of anticancer drugs in dogs and monkeys for the prediction of qualitative toxicities in man

The usefulness of dogs and monkeys in predicting potential qualitative drug toxicity in man was examined retrospectively for twenty-five anticancer compounds of diverse chemical and functional classification. It was found that the large animal screen served to alert the physician to a significant proportion of the total spectrum of drug effects, which were encountered during the clinical use of a new toxic compound. The dog and monkey correctly predicted bone marrow depression, gastrointestinal disturbance, and hepatotoxicity for each drug producing these effects in the clinic; in the case of renal, cardiovascular, and neuromuscular toxicity, however, the large animal screen failed in each instance to predict one drug that produced these toxicities in man. The correct predictions were accomplished at the expense of a high percentage of false positives, which resulted from the necessity of using severely toxic dose levels in order to demonstrate all potential toxicities inherent in any compound. While organ system toxicity observed during an animal study can never be disregarded, it should be viewed with an understanding of certain limitations of animal toxicologic data. While toxicity may develop in man in an organ system predicted susceptible by an animal, a different specific clinical or chemical parameter may be involved. The adverse reaction may appear in man at a greater or lesser dose level or may follow a different order of appearance in relationship to the total spectrum of qualitative toxicity inherent in the compound.

P. S. Schein, R. D. Davis, S. Carter, J. Newman, D. R. Schein, and D. P. Rall
Bethesda, Md.

Laboratory of Toxicology, Cancer Therapy Evaluation Branch, and Program Analysis Branch, National Cancer Institute, National Institutes of Health

One of the most controversial areas in the development of new drugs for use in man involves the efficacy of animal toxicology studies as a predictive system for qualitative toxicity. The general approach of evaluating a new compound for safety in several animal species is performed in an attempt to generate information that

may alert the clinical pharmacologist to potential hazards. The process requires that toxicity data be extrapolated from one species to another, and implicit is the assumption that particular animal species have significant predictive value for toxicity in man and that important toxicity will not go unpredicted. While this topic has

Table I

<i>Drug</i>	<i>Chemical classification</i>	<i>Mechanism of action</i>	<i>Route</i>	<i>Clinical MTD</i>	<i>Principal clinical toxicities</i>	<i>Dog</i>	<i>Monkey</i>
NSC-13875 Melamine, hexamethyl Ref. 13, 64, 111	Triazene	Alkylating agent	PO	12 mg./Kg./day \times 21 (480 mg./M. ²) 8 mg./Kg./day \times 90 (320 mg./M. ²)	Leukopenia Thrombocytopenia Vomiting	+	+
NSC-17256E Pregn-4-ene-3, 11, 20-trione, 6 α -methyl- Ref. 55	Steroid	Steroid hormone	PO	9 mg./Kg./day \times 56 (360 mg./M. ²)	Diarrhea Mild leukopenia	+	-
NSC-19893 Uracil, 5-fluoro- Ref. 22, 94, 102	Pyrimidine	Antimetabolite	IV	15 mg./Kg./day \times 5 (600 mg./M. ²) 7.5 mg./Kg. q.o.d.) (300 mg./M. ²)	Diarrhea Bone marrow depression Stomatitis	+	+
NSC-24559 Mithramycin (USAN) Ref. 21, 59, 84, 88, 97	Tricyclic glycoside	Antibiotic, complexes with DNA, inhibits RNA synthesis	IV	0.025 mg./Kg./day \times 8-10 (1 mg./M. ²)	Bleeding with or with- out thrombocytopenia Bone marrow depression Hypocalcemia Vomiting	-	-
NSC-2698-0 Mitomycin C Ref. 3, 31, 32, 54, 71, 103, 105	Indolequinone	Antibiotic, alkylating agent	IV	0.05 mg./Kg./day \times 5-6 (2 mg./M. ²)	Bone marrow depression Vomiting	+	+
NSC-40774 9H-Purine, 6-(methylthio)- 9- β -D-ribofuranosyl-, dihydrate Ref. 7, 49, 68	Purine nucleoside	Antimetabolite	IV	5 mg./Kg./day \times 5 (200 mg./M. ²)	Stomatitis Mucositis Leukopenia	-	-
NSC-45388 Imidazole-4 (or 5)-carboxam- ide, 5 (or 4)-(3, 3-di- methyl-1-triazeno)- Ref. 64, 67, 95	Triazeno imida- zole	Unknown	IV	5 mg./Kg./day \times 10 (200 mg./M. ²)	Vomiting Bone marrow depression Hepatotoxicity	-	-

NSC-2698-0 Mitomycin C Ref. 3, 31, 32, 54, 71, 103, 105	Indolequinone	Antibiotic, alkylating agent	IV	0.05 mg./Kg./day × 5-6 (2 mg./M. ²)	Bone marrow depression Vomiting	+	+
NSC-40774 9H-Purine, 6-(methylthio)- 9-β-D-ribofuranosyl-, dihydrate Ref. 7, 49, 68	Purine nucleoside	Antimetabolite	IV	5 mg./Kg./day × 5 (200 mg./M. ²)	Stomatitis Mucositis Leukopenia	-	-
NSC-45388 Imidazole-4 (or 5)-carboxam- ide, 5 (or 4)-(3, 3-di- methyl-1-triazeno)- Ref. 64, 67, 95	Triazeno imida- zole	Unknown	IV	5 mg./Kg./day × 10 (200 mg./M. ²)	Vomiting Bone marrow depression Hepatotoxicity	-	-
NSC-51095 Ammonium, trimethyl purine- 6-yl-chloride Ref. 50, 101	Purine	Antimetabolite	IV	30 mg./Kg./day × 7 (1,200 mg./M. ²)	Cholinergic stimulation Bone marrow depression Neurological	+	+
NSC-52947 Pactamycin Ref. 5, 110	Unknown antibi- otic	Antibiotic	IV	0.083 mg./Kg./day × 7 (3.32 mg./M. ²)	Cardiopulmonary Vomiting Bone marrow depression	-	-
NSC-53398 Restrictocin	Polypeptide	Antibiotic	IV	0.008 mg./Kg./day × 5 (0.32 mg./M. ²)	Hypotension Neurologic	-	-
NSC-56408 7 H-Pyrrolo [2, 3-d] pyrimidine, 4-amino-7- β-D-ribofuranosyl-(Tuber- cidin) Ref. 1, 6	Deazapurine nu- cleoside	Antibiotic, antimetabo- lite	IV	1.5 mg./Kg./week × 2 then 0.75 mg./Kg./ month (60 mg./M. ² /→30 mg./M. ²)	Injection-site Vomiting Bone marrow depression Hepatotoxicity	+	+
NSC-56410 Porfiromycin Ref. 24, 33, 40, 66, 106	Indole quinone	Antibiotic, alkylating agent	IV	0.6 mg./Kg./week × 4-6 (24.0 mg./M. ²) 0.5 mg./Kg. twice a week × 3-4 (20.0 mg./M. ²) 0.2 mg./Kg./day × 21 (8.0 mg./M. ²)	Bone marrow depression Vomiting	+	+

Continued on next page.

Table I. *Cont'd*

Drug	Chemical classification	Mechanism of action	Route	Clinical MTD	Principal clinical toxicities	Dog	Monkey
NSC-56654 Azotomycin (USAN) Ref. 2, 105	Diazonium alkylating agent	Antibiotic, glutamine antagonist	IV	2 mg./Kg./day × 8-10 (80 mg./M. ²)	Bone marrow depression Vomiting	+	- No data
NSC-58404 Glycine, <i>N</i> -(diazooacetyl), hydrazide Ref. 38	Diazo hydrazide	Unknown	IV	37 mg./Kg./day × 21 (1.48 Gm./M. ²)	Vomiting	+	-
NSC-60339 Terephthalanilide, 2-chloro-4', 4''-di-2-imidazolin-2-yl- Ref. 8, 48, 78, 93	Terephthalanilide	Unknown	IV	6 mg./Kg./day × 10 (240 mg./M. ²) 6 mg./Kg. twice a week × 8 (240 mg./M. ²)	Ophthalmoplegia Abdominal cramps	- -	+ -
NSC-60387 Tylocerebrine	Polycyclic alkaloid	Antibiotic	IV	0.40 mg./Kg./day × 24 (16 mg./M. ²)	Neurologic Vomiting Bone marrow depression	- + +	- + +
NSC-62512 Acetophenone, 2-(dimethylamino)-3', 4'-dihydroxy-, hydrochloride	Acetophenone	Unknown	IV	275 mg./Kg./day × 10 (10.0 Gm./M. ²)	Bone marrow depression Vomiting Hepatotoxicity	- + +	- + +
NSC-63878 Cytosine, 1-β-D arabinofuranosyl-, monohydrochloride Ref. 16, 20, 46, 58, 65, 81, 98	Pyrimidine nucleoside	Antimetabolite	IV	3.5 mg./Kg./day × 10 (IV push) (140 mg./M. ²) 1-3 mg./Kg./day (24 hr. infus. to toxicity) (40-120 mg./M. ²)	Bone marrow depression Vomiting Hepatotoxicity Stomatitis	+	+
NSC-65346 Sangivamycin Ref. 19	Deazapurine nucleoside	Antibiotic, antimetabolite	IV	Not fully established	Cardiac Bone marrow depression	- +	- -
NSC-68626 Hydrazine, 1-acetyl-2-picolinoyl-	Hydrazine	Chelating agent, peroxide former	IV	12-15 mg./Kg./day × 7-10 (480-600 mg./M. ²)	Neurologic	+	+

NSC-63878 Cytosine, 1- β -D arabinofur- anosyl-, monohydrochloride Ref. 16, 20, 46, 58, 65, 81, 98	Pyrimidine nu- cleoside	Antimetabonite	IV	0.5 mg./Kg./day \times 10 (IV push) (140 mg./M. ²) 1-3 mg./Kg./day (24 hr. infus. to toxicity) (40-120 mg./M. ²)	Vomiting Hepatotoxicity Stomatitis	+	-	-
NSC-65346 Sangivamycin Ref. 19	Deazapurine nucleoside	Antibiotic, antimetabo- lite	IV	Not fully established	Cardiac Bone marrow depression	-	-	+
NSC-68626 Hydrazine, 1-acetyl-2-pico- linoyl-	Hydrazine	Chelating agent, perox- ide former	IV	12-15 mg./Kg./day \times 7-10 (480-600 mg./M. ²)	Neurologic	+	+	
NSC-69945 Phosphorodiamidic acid, N,N-bis (2-chloroethyl)-, compd. with cyclohexyl- amine (1:1) Ref. 70, 76	Phosphorus-con- taining alkylat- ing agent	Alkylating agent	IV	1.5 mg./Kg./day \times 9 (60 mg./M. ²) 2.5 mg./Kg. twice a week \times 2 (100 mg./M. ²) 5.0 mg./Kg./week \times 19 (200 mg./M. ²)	Bone marrow depression Vomiting Hemorrhagic cystitis	+	+	+
NSC-82151 Daunomycin Ref. 18, 29, 99, 100	Tetracyclic gly- coside, poly- hydroxyanthra- quinone glyco- side	Antibiotic, complexes with DNA, inhibits RNA synthesis	IV	1 mg./Kg./day \times 3-5 (45 mg./M. ²)	Bone marrow depression Cardiac	+	+	-
NSC-85998 Streptozotocin Ref. 86, 92	Nitrosourea amino sugar	Antibiotic, depression of NAD	IV	25-75 mg./Kg./week (1-3 Gm./M. ²) >40 mg./Kg./day \times 5 (>1,600 mg./M. ²)	Injection site Vomiting	-	-	+
NSC-94100 Mannitol, 1,6-dibromo-1,6- dideoxy-, D- Ref. 17, 30, 52	Halogenated re- duced sugar	Alkylating agent	PO	6 mg./Kg. to toxicity or therapeutic effect (240 mg./M. ²)	Bone marrow depression	+	+	
NSC-409, 962 Urea, 1,3-bis(2-chloroethyl)- 1-nitroso- Ref. 26, 27, 28, 36, 37, 53, 77, 85, 87, 89, 108	Nitrosourea	Alkylating agent		2.5 mg./Kg./day \times 3 (100 mg./M. ²) 7.5 mg./Kg./day \times 1 (300 mg./M. ²)	Bone marrow depression Hepatotoxicity Vomiting	+	+	+

Table II. Number of animals and patients receiving each drug

Drug	Route	Dog	Monkey	Man
NSC-13875 Melamine, hexamethyl-	PO	Beagle—10	<i>Macaca mulatta</i> —3	100
NSC-17256E Pregn-4-ene-3,11,20-trione,6 α -methyl-	PO	Beagle—4	<i>Macaca mulatta</i> —2	93
NSC-19893 Uracil, 5-fluoro-	IV	Beagle—15	<i>Macaca mulatta</i> —3	> 100
NSC-24559 Mithramycin (USAN)	IV	Beagle—14	<i>Macaca irus</i> —6	900
NSC-26980 Mitomycin C	IV	Mongrel—28 Beagle—9	<i>Macaca mulatta</i> —6	1,004
NSC-40774 9H-Purine,6-(methylthio)-9- β -D-ribo- furanosyl-, dihydrate	IV	Beagle—9	<i>Macaca mulatta</i> —7	70
NSC-45388 Imidazole-4(or 5)-carboxamide, 5(or 4)-(3,3-dimethyl-1-triazeno)-	IV	Beagle—14	<i>Macaca mulatta</i> —7	250
NSC-51095 Ammonium, trimethylpurin-6-yl-chloride	IV	Beagle—13	<i>Macaca mulatta</i> —3	36
NSC-52947 Pactamycin	IV	Beagle—18	<i>Macaca mulatta</i> —6	81
NSC-53398 Restrictocin	IV	Beagle—12	<i>Macaca mulatta</i> —7 <i>Macaca speciosa</i> —4	12
NSC-56408 Tubercidin	IV	Beagle—20	<i>Macaca mulatta</i> —3	132
NSC-56410 Porfiromycin	IV	Beagle—9	—	93
NSC-56654 Azotomycin (USAN)	IV	Beagle—15	—	105
NSC-58404 Glycine, N-(diazooacetyl)-, hydrazide	IV	Mongrel—10 Beagle—6	<i>Macaca irus</i> —7	22
NSC-60339 Terephthalanilide, 2-chloro-4',4''-di-2- imidazolin-2-yl	IV	Beagle—4	<i>Macaca mulatta</i> —6	35
NSC-60387 Tylocerebrine	IV	Mongrel—5 Beagle—7	<i>Macaca irus</i> —7	10
NSC-62512 Acetophenone, 2-(dimethylamino)-3', 4'-dihydroxy-, hydrochloride	IV	Beagle—18	<i>Macaca mulatta</i> —4	59
NSC-63878 Cytosine, 1- β -D-arabinofuranosyl-, monohydrochloride	IV	Mongrel—10	<i>Macaca mulatta</i> —20	> 100
NSC-65346 Sangivamycin	IV	Mongrel—16	<i>Macaca mulatta</i> —4	45
NSC-68626 Hydrazine, 1-acetyl-2-picolinoyl-	IV	Beagle—17	<i>Macaca irus</i> —8	40

Table

NSC-68
Phosph
chlor
hexyl
NSC-82
Daunor
NSC-85
Strepto
NSC-94
Mannit
NSC-40
Urea, 1

been t
and d
studies
are fou
In this
tailed s
twenty-
key, and

Meth

Twen
chemica
examine
there ha
availabl
cies and
toxicolog
under
Chemotl
Protocol
the Nat
monkeys
were us
between
munized
and pu
weighed
were scr
of parasi
Acute let
and suba
carried o

Table II. *Cont'd*

Monkey	Man	Drug	Route	Dog	Monkey	Man
atta-3	100	NSC-69945	IV	Beagle-14	<i>Macaca mulatta</i> -7	55
atta-2	93	Phosphorodiamidic acid, <i>N,N</i> -bis (2-chloroethyl)-, compd. with cyclohexylamine (1:1)				
latta-3	> 100	NSC-82151	IV	Beagle-18	<i>Cercopithecus aethiops</i> -5	> 100
atta-6	900	Daunomycin				
latta-6	1,004	NSC-85998	IV	Beagle-12	<i>Macaca mulatta</i> -6	5
latta-7	70	Streptozotocin				
latta-7	250	NSC-94100	PO	Beagle-9	<i>Macaca mulatta</i> -7	50
latta-3	36	Mannitol, 1,6-dibromo-1,6-dideoxy-, <i>D</i> -				
latta-6	81	NSC-409,962	IV	Beagle-47	<i>Macaca mulatta</i> -15	> 200
latta-7	12	Urea, 1,3-bis(2-chloroethyl)-1-nitroso-				
latta-3	132					
—	93					
—	105					
us-7	22					
ulatta-6	35					
us-7	10					
ulatta-4	59					
ulatta-20	> 100					
ulatta-4	45					
rus-8	40					

been the subject of numerous editorials and discussions, only two comparative studies with the use of relatively few drugs are found in the pharmacologic literature. In this communication, we present a detailed study of the qualitative toxicity of twenty-five anticancer drugs in dog, monkey, and man.

Methods

Twenty-five compounds of diverse chemical and functional classification are examined. For a compound to be included, there had to be detailed toxicologic data available for at least one large animal species and for man (Table I). The animal toxicologic evaluations were performed under the guidelines of The Cancer Chemotherapy National Service Center Protocol¹⁵ at laboratories under contract to the National Cancer Institute. Dogs and monkeys (for species used see Table II) were used exclusively. The dogs weighed between 7 to 9 kilograms and were immunized against distemper and hepatitis and purged of parasites. The monkeys weighed between 2 to 5 kilograms and were screened for tuberculosis and purged of parasites prior to being placed on test. Acute lethal toxicity was elucidated in dogs and subacute repeated-dose toxicology was carried out in both dogs and monkeys. The

numbers and types of animals used for each of the twenty-five drugs are listed in Table II. In all cases, doses administered to the animals were in excess of the maximum tolerated dose level (MTD) (that dose which produced only minimal reversible toxicity) as summarized in Tables III and IV. When treatment ended, half of the animals on each dose level were randomly selected for immediate killing, while the remaining animals were observed for delayed toxicity and reversibility of toxicity. The clinical studies were carried out at the Clinical Center Program of the National Cancer Institute, one or several of the Cancer Chemotherapy Cooperative Study groups, or at an independent institute. The patient data were cumulated from individual patient records, literature publications, and personal communications. Approximately 170 parameters were examined, and a set of criteria of toxicity in animals was established (Table V).

The criteria for hematologic and chemical toxicity in man were based on the range of normal values outlined by the Clinical Pathology Department of the Clinical Center, National Institutes of Health. In patients in whom base-line values were abnormal as a result of primary disease, metastasis, or prior therapy, decisions re-

Table III. Maximum tolerated dose—mg./Kg. (mg./M.²)*

Drug	Route	Dog	Monkey	Man
NSC-13875 Melamine, hexamethyl	PO	<50 mg./Kg./day × 10 (<1,000 mg./M. ²)	21 mg./Kg./day × 15 (252 mg./M. ²)—IV	12 mg./Kg./day × 21 (480 mg./M. ²) 8 mg./Kg./day × 90 (320 mg./M. ²)
NSC-17256E Pregn-4-ene-3,11,20-trione, 6α-methyl-	PO	<30 mg./Kg./day × 28 (<600 mg./M. ²)	<30 mg./Kg./day × 28 (<360 mg./M. ²)	9 mg./Kg./day × 56 (360 mg./M. ²)
NSC-19893 Uracil, 5-fluoro-	IV	5 mg./Kg./day × 5 (100 mg./M. ²) 10 mg./Kg./week × 6 (200 mg./M. ²)	7.5 mg./Kg./day × 6 (90 mg./M. ²)	15 mg./Kg./day × 5 (600 mg./M. ²) 7.5 mg./Kg. q.o.d. (300 mg./M. ²)
NSC-24559 Mithramycin (USAN)	IV	0.025 mg./Kg./day × 24 (0.5 mg./M. ²)	0.025 mg./Kg./day × 24 (0.3 mg./M. ²)	0.025 mg./Kg./day × 5 (1 mg./M. ²)
NSC-26980 Mitomycin C	IV	<0.063 mg./Kg./day × 10 (<1.26 mg./M. ²)	0.2 mg./Kg./day × 10 (2.4 mg./M. ²)	0.05 mg./Kg./day × 5-6 (2 mg./M. ²)
NSC-40774 9H-Purine, 6-(methylthio)-9-β-D-ribo- furanosyl-, dihydrate	IV	4 mg./Kg./day × 10 (80 mg./M. ²)	4 mg./Kg./day × 14 (48 mg./M. ²)	5 mg./Kg./day × 5 (200 mg./M. ²)
NSC-45388 Imidazole-4(or 5)-carboxamide, 5(or 4)-(3,3-dimethyl-1-triazeno)-	IV	2.5 mg./Kg./day × 28 (50 mg./M. ²)	15-30 mg./Kg./day × 28 (185-370 mg./M. ²)	5 mg./Kg./day × 10 (200 mg./M. ²)
NSC-51095 Ammonium, trimethylpurin-6-yl-chloride	IV	8 mg./Kg./day × 28 (160 mg./M. ²)	30 mg./Kg./day × 28 (360 mg./M. ²)—PO	30 mg./Kg./day × 7 (1,200 mg./M. ²)
NSC-52947 Pactamycin	IV	0.02 mg./Kg./day × 28 (0.4 mg./M. ²)	0.02 mg./Kg./day × 23 (0.24 mg./M. ²)	0.083 mg./Kg./day × 27 (3.32 mg./M. ²)
NSC-53398 Restrictocin	IV	<0.01 mg./Kg./day × 28 (<0.2 mg./M. ²)	0.02 mg./Kg./day × 28 (<0.24 mg./M. ²)	0.008 mg./Kg./day × 5 (0.32 mg./M. ²)
NSC-56408 Tubercidin	IV	<0.25 mg./Kg./day × 28 (<0.50 mg./M. ²)	<1 mg./Kg./day × 24 (<12 mg./M. ²)	1.5 mg./Kg./week × 2 then 0.75 mg./Kg./month (60 mg./M. ² → 30 mg./M. ²)

NSC-51095 Ammonium, trimethylpurin-6-yl-chloride	IV	8 mg./Kg./day × 28 (160 mg./M. ²)	30 mg./Kg./day × 28 (360 mg./M. ²)—PO	30 mg./Kg./day × 7 (1,200 mg./M. ²)
NSC-52947 Pactamycin	IV	0.02 mg./Kg./day × 28 (0.4 mg./M. ²)	0.02 mg./Kg./day × 23 (0.24 mg./M. ²)	0.083 mg./Kg./day × 27 (3.32 mg./M. ²)
NSC-53398 Restrictocin	IV	<0.01 mg./Kg./day × 28 (<0.2 mg./M. ²)	0.02 mg./Kg./day × 28 (<0.24 mg./M. ²)	0.008 mg./Kg./day × 5 (0.32 mg./M. ²)
NSC-56408 Tubercidin	IV	<0.25 mg./Kg./day × 28 (<0.50 mg./M. ²)	<1 mg./Kg./day × 24 (<12 mg./M. ²)	1.5 mg./Kg./week × 2 then 0.75 mg./Kg./month (60 mg./M. ² → 30 mg./M. ²)
NSC-56410 Porfiriomycin	IV	<0.310 mg./Kg./day × 10 (<6.2 mg./M. ²)	—	0.6 mg./Kg./week × 4-6 (24.0 mg./M. ²) 0.5 mg./Kg. twice a week × 3-4 (20.0 mg./M. ²) 0.2 mg./Kg./day × 21 (8.0 mg./M. ²)
NSC-56654 Azotomycin (USAN)	IV	0.1 mg./Kg./day × 14 (2.0 mg./M. ²)	—	2 mg./Kg./day × 8-10 (80 mg./M. ²)
NSC-58404 Glycine, N-(diazooacetyl)-, hydrazide	IV	10 mg./Kg./day × 24 (200 mg./M. ²)	20 mg./Kg./day × 24 (240 mg./M. ²)	37 mg./Kg./day × 21 (1.48 Gm./M. ²)
NSC-60339 Terephthalanilide, 2-chloro-4',4''-di-2- imidazolin-2-yl-	IV	0.5 mg./Kg./day × 28 (10 mg./M. ²)	<1 mg./Kg./day × 28 (<12 mg./M. ²)	6 mg./Kg./day × 10 (240 mg./M. ²) 6 mg./Kg. twice a week × 8 (240 mg./M. ²)
NSC-60387 Tylocerebrine	IV	0.125 mg./Kg./day × 24 (2.50 mg./M. ²)	0.25 mg./Kg./day × 24 (3.0 mg./M. ²)	0.40 mg./Kg./day × 24 (16 mg./M. ²)
NSC-62512 Acetophenone, 2-(dimethylamino)-3', 4'-dihydroxy-, hydrochloride	IV	87 mg./Kg./day × 28 (1.74 Gm./M. ²)	260 mg./Kg./day × 28 (3.12 Gm./M. ²)	275 mg./Kg./day × 10 (10.0 Gm./M. ²)
NSC-63878 Cytosine, 1-β-D-arabinofuranosyl-, monohydrochloride	IV	6 mg./Kg./day × 15 (120 mg./M. ²)	12 mg./Kg./day × 15 (144 mg./M. ²)	3.5 mg./Kg./day × 10 (IV push) (140 mg./M. ²) 1-3 mg./Kg./day (24 hr. in- fus. to toxicity) (40-120 mg./M. ²)

*Maximum tolerated dose—the highest dose that either produced no toxicity or that produced toxicity that was in all cases rapidly and completely reversible.

Continued on next page.

Table III. *Cont'd*

Drug	Route	Dog	Monkey	Man
NSC-65346 Sangivamycin	IV	0.050 mg./Kg./day × 24 (1 mg./M. ²)	<0.4 mg./Kg./day × 28 (<4.8 mg./M. ²)	Not fully established
NSC-68626 Hydrazine, 1-acetyl-2-picolinoyl-	IV	12 mg./Kg./day × 24 (240 mg./M. ²)	24 mg./Kg./day × 24 (288 mg./M. ²)	12-15 mg./Kg./day × 7-10 (480-600 mg./M. ²)
NSC-69945 Phosphorodiamidic acid, N,N,-bis (2-chloroethyl)-cpd. with cyclohexylamine (1:1)	IV	4 mg./Kg./day × 14 (80 mg./M. ²)	2 mg./Kg./day × 14 (24 mg./M. ²)	1.5 mg./Kg./day × 9 (60 mg./M. ²) 2.5 mg./Kg. twice a week × 2 (100 mg./M. ²) 5.0 mg./Kg./week × 19 (200 mg./M. ²)
NSC-82151 Daunomycin	IV	<0.5 mg./Kg./day × 12 (<10 mg./M. ²) 2.5 mg./Kg./day × 1 (50 mg./M. ²) <1 mg./Kg./week × 30 (<20 mg./M. ²)	0.25 mg./Kg./day × 14 (3.0 mg./M. ²)	1 mg./Kg./day × 3-5 (45 mg./M. ²)
NSC-85998 Streptozotocin	IV	1 mg./Kg./day × 14 (20 mg./M. ²)	3 mg./Kg./day × 14 (36 mg./M. ²)	>40 mg./Kg./day × 5 (>1,600 mg./M. ²)
NSC-94100 Mannitol, 1,6-dibromo-1,6-dideoxy-,D	PO	13 mg./Kg./day × 90 (260 mg./M. ²) 25 mg./Kg./day × 14 (500 mg./M. ²)	10 mg./Kg./day × 200 (120 mg./M. ²)	6 mg./Kg. to toxicity or therapeutic effect (240 mg./M. ²)
NSC-409,962 Urea, 1,3-bis(2-chloroethyl)-1-nitroso-	IV	1.25 mg./Kg./day × 15 (25 mg./M. ²)	1.25 mg./Kg./day × 15 (15 mg./M. ²)	2.5 mg./Kg./day × 3 (100 mg./M. ²) 7.5 mg./Kg./day × 1 (300 mg./M. ²)

gar-
on t
enc
ten
T.
each
in t
If a
serv
and
nate
Th
from
1231
com
Th
each
was
ity o
was
syste
in be
anim
was i
true
demo
was)
negat
false)
tive,
negat
incide
categ
recte
eters
sent c
listed
compu
and a
within
the or
for to
was ca
followe
mal sp
lated a
negativ
true pc
false m
toxicity

Mannitol, 1,6-dibromo-1,6-dideoxy-, D	(260 mg./M. ²) 25 mg./Kg./day × 14 (500 mg./M. ²)	(120 mg./M. ²)	therapeutic effect (240 mg./M. ²)
IV	1.25 mg./Kg./day × 15 (25 mg./M. ²)	1.25 mg./Kg./day × 15 (15 mg./M. ²)	2.5 mg./Kg./day × 3 (100 mg./M. ²) 7.5 mg./Kg./day × 1 (300 mg./M. ²)
NSC-409,962			
Urea, 1,3-bis(2-chloroethyl)-1-nitroso-			

garding specificity of toxicity were based on the results of serial testing and consistency with findings in similarly treated patients.

The incidence of a given toxicity for each drug was not given special weight in the evaluation of a specific parameter. If an abnormal symptom or sign was observed in more than one animal or patient and considered drug related, it was designated as positive for toxicity.

The data were transferred to IBM cards from IBM mark sense sheets by an IBM 1231 Optical Mark Sense Reader and then computer processed.

The relative frequency of toxicity for each parameter in the individual species was determined (Table VI), and the ability of the animal system to predict for man was defined with the use of the following system: true positive, toxicity was noted in both animal and man; false positive, animal was positive for toxicity but man was negative (an index of overprediction); true negative, neither animal nor man demonstrated toxicity; false negative, man was positive for toxicity but animal was negative (an index of underprediction); false negative/true positive plus false negative, an analysis of the number of false negative predictions as a per cent of the incidence of toxicity in man. The latter category has been designated the "corrected false negative." Groups of parameters whose composite served to represent different specific organ systems, as listed in Tables XI, XII, and XIII, were computed as follows: If for a given animal and a given drug any individual parameter within the group was found to be positive, the organ system was designated positive for toxicity; otherwise, the organ system was called negative. The same process was followed for man. The results for the animal species were then compared and tabulated as true positive, false positive, true negative, false negative, or false negative/true positive plus false negative (corrected false negative) in regard to predicting for toxicity in man.

Results

Prediction of organ system toxicity

Bone marrow. The combined animal screen predicted for all drugs that produced anemia, but at the expense of 44 per cent overprediction (Table IX). It can be noted that neither species predicted for all instances of lowered red blood cell parameters. The dogs were more accurate and efficient, having predicted a higher percentage of true positive at the expense of less false positive while failing to predict for only one of twelve compounds that produced anemia in man (Table VII). However, it can be seen that the monkey data served to correct for this one false negative prediction.

The monkey and dog performed equally well in predicting for leukopenia, with a 60 or 61 per cent individual and 68 per cent combined true positives (Tables VII, VIII, and IX). However, false negative, or underprediction, which was not corrected by the combination was found for both species. Table X presents those compounds in which one or both animal species failed to predict for leukopenia. In addition, the individual animal data did not distinguish between general depression of the white blood cell count and specific toxicity to the granulocytic versus lymphocytic series; the distinctive granulocytopenia of nine of twenty compounds was not predicted by the animal screen.

Thrombocytopenia was correctly predicted for thirteen of the eighteen compounds that caused this toxicity in man (Table IX). The monkey showed significant underprediction, with 41 per cent false negatives and a corrected false negative index of 56 per cent (Table VIII). The specific compounds for which thrombocytopenia was not predicted are summarized in Table X.

The combination of dog and monkey as a toxicologic screen predicted for the myelosuppression for each of the twenty-two compounds that produced this toxicity in man (Table XIII). By examining Tables XI and XII, it can be seen that the lack

Table IV. Maximum dose administered to each of the three species

Drug	Route	Dog	Monkey	Man
NSC-13875 Melamine, hexamethyl-	PO	260 mg./Kg./day \times 8 (5,200 mg./M. ²)	84 mg./Kg./day \times 15 (1,008 mg./M. ²)—IV	12 mg./Kg./day \times 21 (480 mg./M. ²) 8 mg./Kg./day \times 90 (320 mg./M. ²)
NSC-17256E Pregn-4-ene-3, 11, 20-trione, 6 α -methyl-	PO	30 mg./Kg./day \times 28 (600 mg./M. ²)	30 mg./Kg./day \times 28 (360 mg./M. ²)	9 mg./Kg./day \times 56 (360 mg./M. ²)
NSC-19893 Uracil, 5-fluoro-	IV	10 mg./Kg./day \times 5 (200 mg./M. ²)	30 mg./Kg./day \times 6 (360 mg./M. ²)	20 mg./Kg./day \times 5 (600 mg./M. ²)
NSC-24559 Mithramycin (USAN)	IV	0.1 mg./Kg./day \times 8-10 (2 mg./M. ²)	0.1 mg./Kg./day \times 4-6 (1.2 mg./M. ²)	0.05 mg./Kg./day \times 5-6 (2.0 mg./M. ²)
NSC-26980 Mitomycin C	IV	3.2 mg./Kg./day \times 10 (64 mg./M. ²)	0.2 mg./Kg./day \times 8 (4.8 mg./M. ²)	0.007 mg./Kg./day \times 5 (0.280 mg./M. ²)
NSC-40774 9H-purine, 6-(methylthio)-9- β -D ribo- furanosyl-, dihydrate	IV	12 mg./Kg./day \times 8 (240 mg./M. ²)	8 mg./Kg./day \times 8 (96 mg./M. ²)	—6.25 mg./Kg./day \times 5 (250 mg./M. ²)
NSC-45388 Imidazole-4(or 5)-carboxamide, 5(or 4)-(3,3-dimethyl-1-triazeno)-	IV	20 mg./Kg./day \times 28 (400 mg./M. ²)	60 mg./Kg./day \times 28 (1,200 mg./M. ²)	5.5 mg./Kg./day \times 10 (220 mg./M. ²)
NSC-51095 Ammonium, trimethylpurin-6-yl-chloride	IV	16 mg./Kg./day \times 28 (320 mg./M. ²)	120 mg./Kg./day \times 9 (1,440 mg./M. ²)—PO	100 mg./Kg./day \times 4 (infusion) (4,000 mg./M. ²) —34 mg./Kg./week \times 3 (push) (1,350 mg./M. ²)
NSC-52947 Pactamycin	IV	0.375 mg./Kg./day \times 6 (7.5 mg./M. ²)	0.4 mg./Kg./day \times 3 (4.8 mg./M. ²)	0.083 mg./Kg./day \times 7 (3.3 mg./M. ²)
NSC-53398 Restrictocin	IV	0.08 mg./Kg./day \times 6 (1.6 mg./M. ²)	0.32 mg./Kg./day \times 8 (3.84 mg./M. ²)	0.032 mg./Kg./day \times 3 (1.28 mg./M. ²)
NSC-56408 Tubercidin	IV	8 mg./Kg./day \times 6 (160 mg./M. ²)	8 mg./Kg./day \times 4 (96 mg./M. ²)	0.3 mg./Kg./day \times 7-10 (12 mg./M. ²)
NSC-56410 Porfiromycin	IV	2.5 mg./Kg./day \times 21 (50 mg./M. ²)	—	0.25 mg./Kg./day \times 21 (10 mg./M. ²)

NSC-56654
Azotomycin (USAN)IV 0.6 mg./Kg./day \times 7-8
(12 mg./M.²)

—

5.9 mg./Kg./day \times 5-10
(236 mg./M.²)

Pactamycin		(7.5 mg./M. ²)	(4.8 mg./M. ²)	(3.3 mg./M. ²)
NSC-53398	IV	0.08 mg./Kg./day × 6	0.32 mg./Kg./day × 8	0.032 mg./Kg./day × 3
Restrictocin		(1.6 mg./M. ²)	(3.84 mg./M. ²)	(1.28 mg./M. ²)
NSC-56408	IV	8 mg./Kg./day × 6	8 mg./Kg./day × 4	0.3 mg./Kg./day × 7-10
Tubercidin		(160 mg./M. ²)	(96 mg./M. ²)	(12 mg./M. ²)
NSC-56410	IV	2.5 mg./Kg./day × 21	—	0.25 mg./Kg./day × 21
Porfiromycin		(50 mg./M. ²)		(10 mg./M. ²)
<hr/>				
NSC-56654	IV	0.6 mg./Kg./day × 7-8	—	5.9 mg./Kg./day × 5-10
Azotomycin (USAN)		(12 mg./M. ²)		(236 mg./M. ²)
NSC-58404	IV	40 mg./Kg./day × 16-24	160 mg./Kg./day × 12	~37 mg./Kg./day × 21
Glycine, N-(diazooacetyl)-,hydrazide		(1,600 mg./M. ²)	(1,920 mg./M. ²)	(1,500 mg./M. ²)
NSC-60339	IV	0.5 mg./Kg./day × 28	2.5 mg./Kg./day × 28	6 mg./Kg./day × 10
Terephthalanilide, 2-chloro-4',4''-di-2-imidazolin-2-yl-		(10 mg./M. ²)	(30 mg./M. ²)	(240 mg./M. ²)
NSC-60387	IV	1 mg./Kg./day × 7-24	2.0 mg./Kg./day × 4	0.5 mg./Kg./day × 21
Tylocerebrine		(20 mg./M. ²)	(24 mg./M. ²)	(20 mg./M. ²)
NSC-62512	IV	400 mg./Kg./day × 10-28	400 mg./Kg./day × 4-28	375 mg./Kg./day × 10
Acetophenone, 2-(dimethylamino)-3',4'-dihydroxy-, hydrochloride		(8,000 mg./M. ²)	(4,800 mg./M. ²)	2-8 hr. infusion (15 Gm./M. ²)
NSC-63878	IV	100 mg./Kg./day × 6	100 mg./Kg./day × 15	4 mg./Kg./day × 10
Cytosine, 1-β-D-arabinofuranosyl-, monohydrochloride		(2,000 mg./M. ²)	(1,200 mg./M. ²)	(160 mg./M. ²)
NSC-65346	IV	0.4 mg./Kg./day × 13	1.6 mg./Kg./day × 16	0.35 mg./Kg./day × 7
Sangivamycin		(8.0 mg./M. ²)	(19.2 mg./M. ²)	(114 mg./M. ²)
NSC-68626	IV	48 mg./Kg./day × 9	96 mg./Kg./day × 18	18 mg./Kg./day × 7-10
Hydrazine, 1-acetyl-2-picolinoyl-		(960 mg./M. ²)	(1,052 mg./M. ²)	(720 mg./M. ²)
NSC-69945	IV	32 mg./Kg./day × 5	16 mg./Kg./day × 7	4 mg./Kg./day × 9
Phosphorodiamidic acid, N,N-bis (2-chloroethyl)-, compd. with cyclohexylamine (1:1)		(640 mg./M. ²)	(192 mg./M. ²)	(160 mg./M. ²)
NSC-82151	IV	1 mg./Kg./day × 12	1 mg./Kg./day × 12	~1.5 mg./Kg./day × 5
Daunomycin		(20 mg./M. ²)	(12 mg./M. ²)	(60 mg./M. ²)
NSC-85998	IV	4.2 mg./Kg./day × 14	12 mg./Kg./day × 14	40 mg./Kg./day × 5
Streptozotocin		(50.4 mg./M. ²)	(144 mg./M. ²)	(1,600 mg./M. ²)
NSC-94100	PO	200 mg./Kg./day × 6	150 mg./Kg./day × 8	6 mg./Kg./day to toxicity or response
Mannitol, 1,6-dibromo-1,6-dideoxy-, D		(4,000 mg./M. ²)	(1,800 mg./M. ²)	(240 mg./M. ²)
NSC-409,962	IV	20 mg./Kg./day × 7	10 mg./Kg./day × 7	10 mg./Kg./day × 3
Urea, 1,3-bis(2-chloroethyl)-1-nitroso-		(400 mg./M. ²)	(120 mg./M. ²)	(300 mg./M. ²)

Table V

A. Criteria employed for the definition and semiquantitation of drug toxicity

Hematology	Dogs		Monkey	
	Mild-moderate	Marked	Mild-moderate	Marked
Hct. (%)	< 38 - ≥ 32	= or < 32	< 37 - ≥ 30	= or < 30
Hgb. (Gm./%)	< 13 - ≥ 11	= or < 11	< 12.3 - ≥ 9.5	= or < 9.5
WBC ($\times 10^3/\text{mm}^3$)	< 6 - ≥ 4	= or < 4	< 6 - ≥ 4	= or < 4
Platelets ($\times 10^3/\text{mm}^3$)	< 175 - ≥ 100	= or < 100	< 200 - ≥ 100	= or < 100

B. Criteria for the semiquantitation of toxicity in dogs and monkeys based on changes from normal pretreatment values

Chemistry	Mild-moderate	Marked
FBS	< 2 × and > 120/mg./100 ml.	= or > 2 × normal pretreatment value
BSP	< 2 × and > 8%	= or > 2 × normal pretreatment value
Alk. p'tase	< 2 × and > 7 Sigma units (dog) > 30 Sigma units (monkey)	= or > 2 × normal pretreatment value
SGOT	< 3 × and > 60 Sigma-Frankel units	= or > 3 × normal pretreatment value
SGPT	< 3 × and > 60 Sigma-Frankel units	= or > 3 × normal pretreatment value
BUN	< 2 × and > 20 mg./100 ml.	= or > 2 × normal pretreatment value
Creatinine	< 2 × and > 1.5 mg./100 ml.	= or > 2 × normal pretreatment value
Bilirubin	< 2 × and > 1.0 mg./100 ml.	= or > 2 × normal pretreatment value
Prothrombin time	< 2 × and > 10 seconds	= or > 2 × normal pretreatment value

of false negatives, or underprediction, was the result of an overlap in information contributed by using the two species; neither animal species predicted for all toxicity, the dog alone missed two compounds, and the monkey failed to predict for one.

Lymphoid organs. The large animal screen overpredicted 76 per cent for lymphopenia and lymph node toxicity. This form of toxicity was noted with only one compound, daunomycin, during the clinical studies in man.

Gastrointestinal tract. The most consistent clinical gastrointestinal toxicity induced by the drugs was vomiting, which appeared with eighteen of twenty-five compounds. The combined screen gave 72 per cent true positives, but missed 12 per cent drug-related emesis (Table IX). Again these results were primarily contributed by the dog, since the monkey gave a significant 57 per cent underprediction with a corrected false negative index of 68 per cent (Table VIII). The same situation holds

true for diarrhea; the dogs gave 36 per cent positives, failing to predict for only one compound, while the monkey failed for six drugs, giving a 67 per cent corrected false negative prediction for that species. The toxicities of stomatitis and gastrointestinal bleeding were observed with relatively few compounds and were generally underpredicted by the animal screen.

The combined dog and monkey screen gave 92 per cent true positives and no false negatives for gastrointestinal toxicity when compared with that of man (Table XIII). When the individual species are compared, it can be seen that the high degree of correspondence is entirely contributed by the dog; since the monkey failed to predict for four of twenty-three compounds (Tables XI and XII).

Liver. The large animal screen predicted for all instances of hepatotoxicity, but at the expense of a 48 per cent overprediction. The comparison of the dog and monkey data failed to disclose a species selectivity.

The individual parameters in the animal

Ta
to:

Bo:

H

H

R

R

N

W

G

L

Ti

Bo

Ly:

L

L

Sf

I

Ga:

W

N:

V

H

Sa

Al

Di

Co

M

H

St

M

Ul

St

Es

St

Sn

P

La

P

Hej

BS

AI

SC

SC

Bil

All

Gl

Pr

Li

Ga

o.

Table VI. Groups of parameters selected as representatives of organ-specific toxicity and their frequency in each of the three species

Drug toxicity		Dog			Monkey			Man		
Monkey										
Rate	Marked									
> 30	= or < 30									
≥ 9.5	= or < 9.5									
> 4	= or < 4									
> 100	= or < 100									
ys based										
Marked										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										
mal pretreatment value										

Table VI. *Cont'd*

	Dog	Monkey	Man
<i>Respiratory</i>			
Dyspnea	7/25	5/23	3/25
Tachypnea	5/25	0/23	1/25
Cough	2/25	0/23	1/25
Bronchospasm	0/25	0/23	1/25
Hemoptysis	0/25	0/23	0/25
Lung pathology	18/25	11/23	0/2
<i>Injection site</i>			
Pain with injection	1/23	0/20	5/23
Inflammation	9/23	8/21	3/23
Thrombosis	3/23	4/21	4/23
Injection site pathology	8/23	7/23	0/0
<i>Integument</i>			
Desquamation	0/25	0/23	1/25
Dermatitis	0/25	1/23	2/25
Loss of pigmentation	0/25	0/23	0/25
Petechiae	0/25	1/23	5/25
Ecchymosis	0/25	2/23	5/25
Cyanosis	2/25	1/23	1/25
Jaundice	3/25	0/23	2/25
Alopecia	1/25	0/23	2/25
<i>Eyes</i>			
Conjunctivitis	2/25	0/23	0/25
Corneal scarring	0/25	0/23	0/25
Cataracts	0/25	0/23	0/25

species were analyzed for abnormalities of the same individual parameters in man (Tables VII, VIII, and IX). In general, the transaminase and alkaline phosphatase proved to be the most useful parameters of liver function. The transaminases, serum glutamic oxaloacetic (SGOT) and serum glutamic pyruvic (SGPT), had the highest true positive index, 23 to 35 per cent; the highest total true prediction of 46 to 55 per cent (true positive plus true negative) and the lowest false negative prediction 0 to 5 per cent with a corrected false negative index of 12.5 per cent (Table IX). These were closely followed by the alkaline phosphatase in the dog which gave a 25 per cent true positive index, a 46 per cent total true prediction, with a 14 per cent corrected false positive prediction for abnormal alkaline phosphatase values in man (Table VII). The bromsulphalein (BSP)

determination gave only an 11 and 32 per cent true positive and total true predictions, respectively, while overpredicting 63 per cent and missing one of three compounds that produced this toxicity in man. The over-all true prediction of bilirubinaemia was 72 per cent with no false negatives. The three instances of prolonged prothrombin time in man were missed by the large animal screen.

Each of the individual liver parameters was then separately analyzed to determine which best predicted for liver toxicity, as represented by the entire group of hepatic function tests (i.e., BSP versus liver, alkaline phosphatase versus liver). The data for each species and the combination are presented in Table XIV. One notable feature of the data was the poor performance of the alkaline phosphatase in monkeys, having failed to predict for seven of ten hepatotoxic compounds. With the use of the monkey and dog combination, the best predictive index was presented by the transaminase determinations which gave a total true prediction of 53 to 55 per cent with a false negative prediction of 0 to 5 per cent while overpredicting for 40 to 46 per cent of compounds tested. The BSP determination gave a 44 to 50 per cent overprediction with the combination of the two species with a total false prediction (false positives plus false negatives) of 55.5 to 66.7 per cent, having failed to predict for four of nine compounds. The data were analyzed to determine whether the alkaline phosphatase and transaminase tests could substitute for all cases in which the BSP in either species gave a true positive and could correct for all false negative predictions. In Table XV it can be seen that the alkaline phosphatase in the dog could substitute for BSP in the five cases in which the latter determination gave a true positive. The addition of the SGPT and SGOT in dog and monkey to the alkaline phosphatase gives added confidence. The data were similarly analyzed to determine whether the alkaline phosphatase and transaminase tests could

only an 11 and 32
and total true pre-
while overpredicting
ng one of three com-
l this toxicity in man.
diction of bilirubine-
with no false nega-
tances of prolonged
man were missed by
n.

dual liver parameters
nalyzed to determine
for liver toxicity, as
tire group of hepatic
SP versus liver, alka-
sus liver). The data
the combination are
IV. One notable fea-
the poor performance
phatase in monkeys,
dict for seven of ten
ds. With the use of
combination, the best
s presented by the
inations which gave
1 of 53 to 55 per cent
prediction of 0 to 5
predicting for 40 to
pounds tested. The
ve a 44 to 50 per cent
he combination of the
total false prediction
false negatives) of
nt, having failed to
nine compounds. The
to determine whether
ase and transaminase
for all cases in which
species gave a true
correct for all false

In Table XV it can
caline phosphatase in
itute for BSP in the
the latter determina-
itive. The addition of
C in dog and monkey
phatase gives added
were similarly ana-
whether the alkaline
nsaminase tests could

Table VII. Dog as a predictor for selected parameter toxicity in man

	TP*	FP†	TN‡	FN§	FN TP + FN
Injection site inflammation	2/22¶ 9.1%	7/22 31.8%	12/22 54.5%	1/22 4.5%	1/3 33.3%
Injection site thrombosis	2/22 9.1%	1/22 4.5%	17/22 77.0%	2/22 9.1%	2/4 50.0%
Dermatitis	0/25	0/25	23/25 92.0%	2/25 8.0%	2/2 100.0%
Petechiae	0/25	0/25	20/25 80.0%	5/25 20.0%	5/5 100.0%
Jaundice	0/25	3/25 12.0%	20/25 80.0%	2/25 8.0%	2/2 100.0%
Flushing	0/25	0/25	21/25 84.0%	4/25 16.0%	4/4 100.0%
Tachycardia	1/25 4.0%	3/25 12.0%	17/25 68.0%	4/25 16.0%	4/5 80.0%
Lethargy	3/25 12.0%	13/25 52.0%	9/25 36.0%	0/25	0/3
Depression of sensorium	3/25 12.0%	7/25 28.0%	14/25 56.0%	1/25 4.0%	1/4 25.0%
Agitation	1/25 4.0%	3/25 12.0%	20/25 80.0%	1/25 4.0%	1/2 50.0%
Ataxia	2/25 8.0%	9/25 36.0%	13/25 52.0%	1/25 4.0%	1/3 33.0%
Tremor	0/25	7/25 28.0%	18/25 72.0%	0/25	0/0
Seizures	1/25 4.0%	4/25 16.0%	20/25 80.0%	0/25	0/1
Dyspnea	1/25	6/25 24.0%	16/25 64.0%	2/25 8.0%	2/3 66.0%
Lacrimation ↑	2/25 8.0%	6/25 24.0%	17/25 68.0%	0/25	0/2
Ophthalmoplegia	0/25	0/25	24/25 96.0%	1/25 4.0%	1/1 100.0%
Weight loss	0/25	23/25 92.0%	2/25 8.0%	0/25	0/0
Vomiting	18/25 72.0%	4/25 16.0%	0/25	3/25 12.0%	3/21 14.4%

*True positive, toxicity was observed both in dogs and in man.

†False positive, toxicity was observed in dogs but not in man.

‡True negative, no toxicity was observed in dogs and man.

§False negative, toxicity was not observed in the dogs but was recorded in man.

|| False negatives

True positives plus false negatives —corrected false negative, an index of false negative prediction which analyzes for only those compounds which produced the specific toxicity in man.

$$\frac{\text{Numerator}}{\text{denominator}} = \frac{\text{the number of drugs producing toxicity}}{\text{the number of drugs tested}}$$

Continued on next page.

Table VII. *Cont'd*

	TP*	FP†	TN‡	FN§	$\frac{FN }{TP + FN}$
Salivation ↑	3/25 12.0%	9/25 36.0%	13/25 52.0%	0/25	0/3
Diarrhea	9/25 36.0%	10/25 40.0%	5/25 20.0%	1/25 4.0%	1/10 100.0%
Melena	0/25	3/25 12.0%	20/25 80.0%	2/25 8.0%	2/2 100.0%
Hematochezia	1/25 4.0%	17/25 68.0%	7/25 28.0%	0/25	0/1
Stomatitis	1/25 4.0%	1/25 4.0%	19/25 76.0%	4/25 16.0%	4/5 80.0%
Hemoglobin ↓	11/25 44.0%	7/25 28.0%	6/25 24.0%	1/25 4.0%	1/12 8.3%
Hematocrit ↓	11/25 44.0%	7/25 28.0%	6/25 24.0%	1/25 4.0%	1/12 8.3%
Red blood cell count ↓	9/25 36.0%	9/25 36.0%	6/25 24.0%	1/25 4.0%	1/10 10.0%
White blood cell count ↓	15/25 60.0%	1/25 4.0%	4/25 16.0%	5/25 20.0%	5/20 25.0%
Thrombocytopenia	13/22 59.1%	3/22 13.6%	3/22 13.6%	3/22 13.6%	3/16 18.7%
Hypercalcemia	0/3	2/3 66.7%	1/3 33.3%	0/3	0/0
BSP retention ↑	2/18 11.1%	12/18 66.6%	4/18 22.2%	0/18	0/2
Alkaline phosphatase ↑	6/24 25.0%	12/24 50.0%	5/24 20.8%	1/24 4.2%	1/7 14.2%
SGOT ↑	5/18 27.8%	7/18 38.9%	4/18 22.2%	2/18 11.1%	2/7 28.4%
SGPT ↑	3/11 27.3%	5/11 45.5%	3/11 27.3%	0/11	0/3
Bilirubinemia	2/16 12.5%	4/16 25.0%	10/16 62.5%	0/16	0/2
Prothrombin time ↑	0/21	4/21 19.0%	15/21 71.4%	2/21 9.5%	2/2 100.0%
Blood urea nitrogen ↑	6/25 24.0%	9/25 36.0%	7/25 28.0%	3/25 12.0%	3/9 33.0%
Creatinine ↑	0/6	1/6 16.7%	5/6 83.3%	0/6	0/0
Hyperglycemia	0/25	4/25 16.0%	20/25 80.0%	1/25 4.0%	1/1 100.0%
Hypoglycemia	0/25	3/25 12.0%	22/25 88.0%	0/25	0/0

Table VII. *Cont'd*

FN§	FN TP + FN		TP*	FP†	TN‡	FN§	FN TP + FN
0/25	0/3	Proteinuria	1/11 9.1%	8/11 72.7%	2/11 18.2%	0/11	0/1
1/25 4.0%	1/10 100.0%	Glycosuria	0/11	4/11 36.4%	7/11 63.6%	0/11	0/0
2/25 8.0%	2/2 100.0%	Bilirubinuria	0/11	5/11 45.5%	6/11 54.5%	0/11	0/0
0/25	0/1	Hematuria	1/11 9.1%	4/11 36.4%	6/11 54.5%	0/11	0/1
4/25 16.0%	4/5 80.0%	Hyposthenuria	0/14	6/14 42.9%	8/14 57.1%	0/14	0/0
1/25 4.0%	1/12 8.3%						
1/25 4.0%	1/12 8.3%						
1/25 4.0%	1/10 10.0%						
5/25 20.0%	5/20 25.0%						
3/22 13.6%	3/16 18.7%						
0/3	0/0						
0/18	0/2						
1/24 4.2%	1/7 14.2%						
2/18 11.1%	2/7 28.4%						
0/11	0/3						
0/16	0/2						
2/21 9.5%	2/2 100.0%						
3/25 12.0%	3/9 33.0%						
0/6	0/0						
1/25 4.0%	1/1 100.0%						
0/25	0/0						

correct for the underprediction, or false negatives, made by the BSP determination. Table XVI presents the results for the four compounds whose liver toxicity was not predicted by BSP. It can be seen that a true positive correlation is made in all cases with the use of both the alkaline phosphatase and SGOT determination and the two laboratory animal species.

Kidney. The large animal screen gave 36 per cent true positives for renal toxicity but underpredicted for one of twenty-five drugs and gave a 56 per cent overprediction (Table XIII). In general, no specific advantage could be attributed to use of either the dog or monkey. The blood urea nitrogen determination gave 24 per cent true positives, while overpredicting 52 per cent, and a 33 per cent corrected false prediction (Table IX). Serum creatinine was measured in man for only seven compounds, and the animal data overpredicted 43 per cent for toxicity. Proteinuria in the animals gave a gross overprediction of 73 per cent for this abnormality in man, as did hematuria and hyposthenuria. The high incidence of renal function abnormalities in the animals had its counterpart in the renal histopathologic changes which were documented for 64 and 56 per cent of drugs in dogs and monkeys, respectively (Table VI).

Neuromuscular. The combination of dog

and monkey predicted for the neuromuscular abnormalities of six of seven compounds which produced this form of toxicity in man. Neither species presented a distinctly greater predictive value, though the monkey demonstrated a lesser degree of overprediction. The screen predicted for two of the three instances of ataxia and the one drug that produced convulsions. The two drugs that produced excessive lacrimation and the one case of ophthalmoplegia had true positive correlations with the monkey data. There was a large overprediction of specific parameters which contributed to the favorable predictive index for neuromuscular toxicity as a group.

Cardiovascular. The combined large animal screen gave a 36 per cent true positive index, with a total true prediction of 64 per cent, and failed to predict for the cardiovascular toxicity of one of ten compounds. The overlap of monkey and dog data contributed to both the higher incidence of true positive and lower index of false negative when compared with predictions of the individual species.

Respiratory. The screen predicted for the respiratory signs or pathology for four of the five compounds at the expense of a large percentage of false positives. In general, neither the dog nor monkey proved superior in predictive value.

Injection site. The injection site toxicity

Table VIII. Monkey as a predictor for selected parameter toxicity in man

	TP*	FP†	TN‡	FN§	FN
					TP + FN
Injection site inflammation	2/20 10.0%	6/20 30.0%	11/20 55.0%	1/20 5.0%	1/3 33.0%
Injection site thrombosis	1/20 5.0%	3/20 15.0%	14/20 70.0%	2/20 10.0%	2/3 66.0%
Dermatitis	0/23	1/23 4.3%	20/23 87.0%	2/23 8.7%	2/2 100.0%
Petechiae	1/23 4.3%	0/23	18/23 78.3%	4/23 17.4%	4/5 80.0%
Jaundice	0/23	0/23	22/23 95.7%	1/23 4.3%	1/1 100.0%
Flushing	1/22 4.5%	0/22	18/22 81.8%	3/22 13.6%	3/4 75.0%
Tachycardia	0/23	1/23 4.3%	17/23 73.9%	5/23 21.7%	5/5 100.0%
Lethargy	1/23 4.3%	7/23 30.4%	13/23 56.5%	2/23 8.7%	2/3 66.0%
Depression of sensorium	2/23 8.7%	6/23 26.1%	13/23 56.5%	2/23 8.7%	2/4 50.0%
Agitation	0/23	0/23	21/23 91.3%	2/23 8.7%	2/2 100.0%
Ataxia	1/23 4.3%	3/23 13.0%	17/23 73.9%	2/23 8.7%	2/3 66.0%
Tremor	0/23	5/23 21.7%	18/23 78.3%	0/23	0/0
Seizures	1/23 4.3%	1/23 4.3%	21/23 91.3%	0/23	0/1
Dyspnea	0/23	5/23 21.7%	15/23 65.0%	3/23 13.0%	3/3 100.0%
Lacrimation ↑	2/23 8.7%	1/23 4.3%	20/23 87.0%	0/23	0/2
Ophthalmoplegia	1/23 4.3%	0/23	22/23 95.7%	0/23	0/1
Weight loss	0/23	14/23 60.9%	9/23 39.1%	0/23	0/0
Vomiting	6/23 26.1%	3/23 13.0%	1/23 4.3%	13/23 56.5%	13/19 68.4%

*True positive, toxicity was observed both in monkeys and in man.

†False positive, toxicity was observed in monkeys but not in man.

‡True negative, no toxicity was observed in monkeys and man.

§False negative, toxicity was not observed in the monkeys but was recorded in man.

|| False negatives

—corrected false negative, an index of false negative prediction which analyzes for only those compounds which produced the specific toxicity in man.

$$\frac{\text{Numerator}}{\text{Denominator}} = \frac{\text{number of drugs producing toxicity}}{\text{number of drugs tested}}$$

211

FN§	FN	
	TP + FN	
1/20	1/3	
5.0%	33.0%	
2/20	2/3	
10.0%	66.0%	
2/23	2/2	
8.7%	100.0%	
4/23	4/5	
17.4%	80.0%	
1/23	1/1	
4.3%	100.0%	
3/22	3/4	
13.6%	75.0%	
5/23	5/5	
21.7%	100.0%	
2/23	2/3	
8.7%	66.0%	
2/23	2/4	
8.7%	50.0%	
2/23	2/2	
8.7%	100.0%	
2/23	2/3	
8.7%	66.0%	
0/23	0/0	
0/23	0/1	
3/23	3/3	
13.0%	100.0%	
0/23	0/2	
0/23	0/1	
0/23	0/0	
13/23	13/19	
56.5%	68.4%	

prediction which analyzes for

Table VIII. *Cont'd*

	TP*	FP†	TN‡	FN§	FN	
					TP + FN	
Salivation ↑	2/23 8.7%	2/23 8.7%	18/23 78.3%	1/23 4.3%	1/3 33.0%	
Diarrhea	3/23 13.0%	6/23 26.1%	8/23 34.8%	6/23 26.1%	6/9 66.0%	
Melena	0/23	0/23	21/23 91.3%	2/23 8.7%	2/2 100.0%	
Hematochezia	0/23	5/23 21.7%	17/23 73.9%	1/23 4.3%	1/1 100.0%	
Stomatitis	0/23	1/23 4.3%	18/23 78.3%	4/23 17.4%	4/4 100.0%	
Hemoglobin ↓	8/22 36.4%	9/22 40.9%	2/22 9.1%	3/22 13.6%	3/11 27.0%	
Hematocrit ↓	8/23 34.8%	10/23 43.5%	2/23 8.7%	3/23 13.0%	3/11 27.3%	
Red blood cell count ↓	6/22 27.3%	11/22 50.0%	2/22 9.1%	3/22 13.6%	3/9 33.0%	
White blood cell count ↓	14/23 60.9%	3/23 13.0%	2/23 8.7%	4/23 17.4%	4/18 22.2%	
Thrombocytopenia	7/22 31.8%	0/22	6/22 27.3%	9/22 40.9%	9/16 56.3%	
Hypercalcemia	0/1	0/1	1/1 100.0%	0/1	0/0	
BSP retention ↑	1/9 11.1%	6/9 66.7%	1/9 11.1%	1/9 11.1%	1/2 50.0%	
Alkaline phosphatase ↑	3/20 15.0%	2/20 10.0%	11/20 55.0%	4/20 20.0%	4/7 57.0%	
SGOT ↑	6/18 33.3%	4/18 22.2%	6/18 33.3%	2/18 11.1%	2/8 25.0%	
SGPT ↑	2/13 15.4%	4/13 30.8%	6/13 46.2%	1/13 7.7%	1/3 33.0%	
Bilirubinemia	0/14	2/14 14.3%	11/14 78.6%	1/14 7.1%	1/1 100.0%	
Prothrombin time ↑	0/6	1/6 16.7%	4/6 66.7%	1/6 16.7%	1/1 100.0%	
Blood urea nitrogen ↑	4/22 18.2%	10/22 45.5%	4/22 18.2%	4/22 18.2%	4/8 50.0%	
Creatinine ↑	0/5	3/5 60.0%	2/5 40.0%	0/5	0/0	
Hyperglycemia	0/22	7/22 31.8%	14/22 63.6%	1/22 4.5%	1/1 100.0%	
Hypoglycemia	0/22	1/22 4.5%	21/22 95.5%	0/22	0/0	
Proteinuria	0/7	3/7 42.9%	3/7 42.9%	1/7 14.3%	1/1 100.0%	

Continued on next page.

Table VIII. *Cont'd*

	TP ^a	FP [†]	TN [†]	FN [§]	FN
					TP + FN
Glycosuria	0/9	2/9 22.2%	7/9 77.8%	0/9	0/0
Bilirubinuria	0/8	1/8 12.5%	7/8 87.5%	0/8	0/0
Hematuria	0/8	1/8 12.5%	6/8 75.0%	1/8 12.5%	1/1 100.0%
Hyposthenuria	0/9	1/9 11.1%	8/9 88.9%	0/9	0/0

of four of six compounds was predicted by the screen with a 36 per cent overprediction. There was no evidence of an animal species specificity. The animal studies did not allow anticipation of pain of injection with four of five compounds that produced this difficulty.

Integument and dermal manifestations of systemic toxicity. The skin of animals failed to correlate with the dermal lesions noted with therapy in man. The screen missed all cases of dermatitis and jaundice while predicting for one of five drugs that caused cutaneous bleeding.

Eyes. There were no cases of drug-related conjunctivitis, corneal scarring, or cataracts associated with therapy in either the monkey or man, but dogs developed conjunctivitis with two of the twenty-five compounds.

Requirement for the administration of highly toxic dose levels for the prediction of specific abnormalities. Moderately to severely toxic dose levels were required in the dog and monkey to disclose the drug-produced thrombocytopenia of mithramycin, mitomycin C, and dibromomannitol and for the leukopenia caused by 5-fluorouracil, the ammonium compound, azotomycin, and sangivamycin. Toxic doses were required for uncovering the hepatotoxicity for the acetophenone compound and cytosine arabinoside and the neurologic toxicity of restrictocin and 1-acetyl-2-picolinoyl-hydrazine.

Discussion

The medical community has taken an ambivalent or generally negative view of the ability of animal toxicology studies to predict for toxic drug effects in man.^{4, 12, 14, 23, 39, 60-62, 96, 113} There has been much support for the concept that prediction from animal data is perilous and that only through trial by ordeal in man can clinical toxicities be delineated. The field of chemical pharmacology has generated a wealth of data clearly demonstrating that species' differences in drug metabolism are the rule rather than the exception, while absorption, distribution, storage, and excretion tend to be similar.⁹⁻¹² However, it must be noted that to the greatest extent these studies were conducted to explain quantitative differences in principal therapeutic effect on the primary target organ (i.e., the length of barbiturate anesthesia) and have been much less concerned or correlated with accompanying qualitative toxicities in other organ systems. Another source of concern regarding the usefulness of animal toxicology stems from examples in the literature in which an animal screen failed to predict or overpredicted for a specific drug effect in man. Several of these reports have been repeatedly cited as classic illustrations of the futility of animal experimentation. Many of these examples should be re-examined in a positive sense for clues that may lead to more realistic expectations from animal data

Table IX. The combination of the dog and monkey as a predictor for selected parameter toxicity in man

FN§	FN
TP + FN	
0/9	0/0
0/8	0/0
1/8	1/1
12.5%	100.0%
0/9	0/0

	TP*	FP†	TN‡	FN§	FN
					TP + FN
Injection site inflammation	2/22¶ 9.1%	10/22 45.5%	9/22 40.9%	1/22 4.5%	1/3 33.0%
Injection site thrombosis	2/22 9.1%	3/22 13.6%	15/22 68.2%	2/22 9.1%	2/4 50.0%
Dermatitis	0/25	1/25 4.0%	22/25 88.0%	2/25 8.0%	2/2 100.0%
Petechiae	1/25 4.0%	0/25	20/25 80.0%	4/25 16.0%	4/5 80.0%
Jaundice	0/25	3/25 12.0%	20/25 80.0%	2/25 8.0%	2/2 100.0%
Flushing	1/25 4.0%	0/25	21/25 84.0%	3/25 12.0%	3/4 75.0%
Tachycardia	1/25 4.0%	3/25 12.0%	17/25 68.0%	4/25 16.0%	4/5 80.0%
Lethargy	3/25 12.0%	15/25 60.0%	7/25 28.0%	0/25	0/3
Depression of sensorium	4/25 16.0%	11/25 44.0%	10/25 40.0%	0/25	0/4
Agitation	1/25 4.0%	3/25 12.0%	20/25 80.0%	1/25 4.0%	1/2 50.0%
Ataxia	2/25 8.0%	11/25 44.0%	11/25 44.0%	1/25 4.0%	1/3 33.0%
Tremor	0/25	9/25 36.0%	16/25 64.0%	0/25	0/0
Seizures	1/25 4.0%	4/25 16.0%	20/25 80.0%	0/25	0/1
Dyspnea	1/25 4.0%	7/25 28.0%	15/25 60.0%	2/25 8.0%	2/3 66.0%
Lacrimation ↑	2/25 8.0%	6/25 24.0%	17/25 68.0%	0/25	0/2
Ophthalmoplegia	1/25 4.0%	0/25	24/25 96.0%	0/25	0/1
Weight loss	0/25	23/25 92.0%	2/25 8.0%	0/25	0/0

*True positive, toxicity was observed in both animals (dog or monkey) and in man.

†False positive, toxicity was observed in the animals but not in man.

‡True negative, no toxicity was observed in the animals and man.

§False negative, toxicity was not observed in the animals but was recorded in man.

|| False negatives

¶ True positives plus false negatives —corrected false negative, an index of false negative prediction which analyzes for only those compounds which produced the specific toxicity in man.

$$\frac{\text{Numerator}}{\text{Denominator}} = \frac{\text{number of drugs producing toxicity}}{\text{number of drugs tested}}$$

Continued on next page.

immunity has taken an rally negative view of l toxicology studies to ug effects in man.^{4, 12,} There has been much oncept that prediction perilous and that only deal in man can clini- lineated. The field of logy has generated a rly demonstrating that in drug metabolism are n the exception, while tion, storage, and ex- similar.⁹⁻¹² However, it to the greatest extent conducted to explain ces in principal thera- e primary target organ (barbiturate anesthesia) uch less concerned or xompaning qualitative organ systems. Another egarding the usefulness y stems from examples which an animal screen or overpredicted for a t in man. Several of been repeatedly cited ns of the futility of ani- n. Many of these exam- examined in a positive at may lead to more ns from animal data

Table IX. *Cont'd*

	TP*	FP†	TN‡	FN§	FN
					TP + FN
Vomiting	18/25 72.0%	4/25 16.0%	0/25	3/25 12.0%	3/21 14.0%
Salivation ↑	3/25 12.0%	9/25 36.0%	13/25 52.0%	0/25	0/3
Diarrhea	9/25 36.0%	11/25 44.0%	4/25 16.0%	1/25 4.0%	1/10 10.0%
Melena	0/25	3/25 12.0%	20/25 80.0%	2/25 8.0%	2/2 100.0%
Hematochezia	1/25 4.0%	18/25 72.0%	6/25 24.0%	0/25	0/1
Stomatitis	1/25 4.0%	2/25 8.0%	18/25 72.0%	4/25 16.0%	4/5 80.0%
Hemoglobin ↓	12/25 48.0%	10/25 40.0%	3/25 12.0%	0/25	0/12
Hematocrit ↓	12/25 48.0%	11/25 44.0%	2/25 8.0%	0/25	0/12
Red blood cell count ↓	10/25 40.0%	12/25 48.0%	3/25 12.0%	0/25	0/10
White blood cell count ↓	17/25 68.0%	3/25 12.0%	2/25 8.0%	3/25 12.0%	3/20 15.0%
Thrombocytopenia	13/24 54.2%	3/24 12.5%	3/24 12.5%	5/24 20.8%	5/18 27.8%
Hypercalcemia	0/3	2/3 66.7%	1/3 33.3%	0/3	0/0
BSP retention ↑	2/19 10.5%	12/19 63.2%	4/19 21.1%	1/19 5.3%	1/3 33.0%
Alkaline phosphatase ↑	6/25 24.0%	12/25 48.0%	5/25 20.0%	2/25 8.0%	2/8 25.0%
SGOT ↑	7/20 35.0%	8/20 40.0%	4/20 20.0%	1/20 5.0%	1/8 12.5%
SGPT ↑	3/13 23.1%	7/13 53.8%	3/13 23.1%	0/13	0/3
Bilirubinemia	2/18 11.1%	5/18 27.8%	11/18 61.1%	0/18	0/2
Prothrombin time ↑	0/21	5/21 23.8%	14/21 66.7%	2/21 9.5%	2/2 100.0%
Blood urea nitrogen ↑	6/25 24.0%	13/25 52.0%	3/25 12.0%	3/25 12.0%	3/9 33.0%
Creatinine ↑	0/7	3/7 42.9%	4/7 57.1%	0/7	0/0
Hyperglycemia	0/25	10/25 40.0%	14/25 56.0%	1/25 4.0%	1/1 100.0%

Table IX. *Cont'd*

FN§	FN TP + FN
3/25 12.0%	3/21 14.0%
0/25	0/3
1/25 4.0%	1/10 10.0%
2/25 8.0%	2/2 100.0%
0/25	0/1
4/25 16.0%	4/5 80.0%
0/25	0/12
0/25	0/12
0/25	0/10
3/25 12.0%	3/20 15.0%
5/24 20.8%	5/18 27.8%
0/3	0/0
1/19 5.3%	1/3 33.0%
2/25 8.0%	2/8 25.0%
1/20 5.0%	1/8 12.5%
0/13	0/3
0/18	0/2
2/21 9.5%	2/2 100.0%
3/25 12.0%	3/9 33.0%
0/7	0/0
1/25 4.0%	1/1 100.0%

	TP*	FP†	TN‡	FN§	FN TP + FN
Hypoglycemia	0/25	4/25 16.0%	21/25 84.0%	0/25	0/0
Proteinuria	1/11 9.1%	8/11 72.7%	2/11 18.2%	0/11	0/1
Glycosuria	0/13	5/13 38.5%	8/13 61.5%	0/13	0/0
Bilirubinuria	0/12	5/12 41.7%	7/12 58.3%	0/12	0/0
Hematuria	1/12 8.2%	5/12 41.7%	6/12 50.0%	0/12	0/1
Hypothenuria	0/15	6/15 40.0%	9/15 60.0%	0/15	0/0

and a better understanding of their limitations. Typical is the case of 6-azauracil, an antimetabolite selected for therapeutic trial in man on the basis of its antitumor activity in a number of rodent tumor systems. During the clinical studies, it was discovered that the principal dose-limiting toxicity involved the central nervous system, as manifested by lethargy leading to a wide range of sensory, motor, and psychic disturbances associated with electroencephalographic changes.⁹⁶ These profound CNS effects occurred at doses that were considered tolerated in a wide range of animal species. However, ample indications of CNS toxicity were available in the animal data, though very high doses were required. In mice, intraperitoneal injection of doses of 3,000 mg. per kilogram produced deep hypnosis, respiratory depression, and loss of response to painful stimuli. Narcosis was produced in rats at a dose of 1,000 mg. per kilogram.¹⁰⁷ Other studies in rats have demonstrated electroencephalographic patterns similar to those noted in man, with subsequent production of profound ataxia.⁷⁵ Temporary paralysis of lower extremities has been reported in dogs and cats after administration of doses of 1,200 and 1,400 mg. per kilogram,

respectively.¹⁰⁷ The quantitative differences in dose level required to achieve neurologic toxicity in animals as compared to those required in man may relate to differences in metabolism and/or species-related variance of target organ sensitivity.

The analysis of the CNS toxicity of 6-azauracil can be used to make two important points. Very large toxic doses should be administered to animals in order to elicit all the potential qualitative toxicities of a compound. If a particular toxicity is observed during the animal studies, it cannot be disregarded since it may appear in man, though at a much different dose level, or may follow a different order of appearance in relationship to the total spectrum of toxic signs. In our comparative study we have encountered a similar situation in the case of 1-acetyl-2-picolinoyl-hydrazine, NSC-68626. The principal dose-limiting toxicity in man is somnolence leading to coma. Neurologic toxicity was seen in the dog in the form of ataxia and convulsions but required high-dose treatment for elicitation.

Another point of concern is that overprediction by a sensitive animal species may prevent a potentially important drug from ever reaching therapeutic trial in

Table X

	Species that failed to predict for leukopenia in man	Species that failed to predict for thrombocytopenia in man
NSC-13875 Melamine, hexamethyl-	Monkey	Monkey
NSC-17256E Pregn-4-ene-3,11,20-trione,6 α -methyl-	Dog, monkey	—
NSC-19893 Uracil, 5-fluoro-	—	Monkey
NSC-24559 Mithramycin (USAN)	Monkey	Dog, monkey
NSC-26980 Mitomycin C	—	Monkey
NSC-40774 9H-Purine, 6-(methylthio)-9- β -D ribo- furanosyl-, dihydrate	Dog	Monkey (not measured in dog)
NSC-51095 Ammonium, trimethylpurin-6-yl-chloride	Monkey	Monkey
NSC-52947 Pactamycin	—	Monkey
NSC-53398 Restrictocin	—	Dog, monkey
NSC-56408 Tubercidin	Dog, monkey	—
NSC-62512 Acetophenone, 2-(dimethylamino)-3',4'-di- hydroxy-, hydrochloride	—	Dog, monkey
NSC-65346 Sangivamycin	Dog	—
NSC-69945 Phosphorodiamidic acid, N,N-bis (2-chlo- roethyl)-, compd. with cyclohexylamine (1:1)	—	Dog, monkey

man. The model often used for illustration is the high degree of toxicity produced by penicillin in the guinea pig.^{14, 60} The original study by Hamre and associates⁴² showed that the single-dose toxicity of penicillin in guinea pigs was similar to that in mice and rabbits. A species difference arose when repeated-dose toxicity studies were conducted in which, at relatively low dosage, the guinea pigs showed a significant mortality rate. The mechanism of this mortality rate has been the

subject of study for many years. The data suggest that death is not a direct drug effect but is, at least in part, related to the replacement of the normal gram-positive intestinal bacterial flora with an overgrowth or superinfection with gram-negative coliforms.²⁵ Thus, the guinea pig would have predicted for one of the principal clinical problems associated with the use of penicillin in man—superinfection; though in the clinic, overgrowth with *Staphylococcus aureus* is more commonly

Table

Injectio
Integum
Cardiov
Respira
Bone m
Lymphc
Gastroin
Liver
Renal
Neuromu

*True
†False
‡True
§False
|| False
|| true p

Table X

Org
Injectio
Integum
Cardiov
Respirat
Bone m
Lympho
Gastroint
Liver
Renal
Neuromu

*True posi
†False pos.
‡True neg.
§False neg.
|| False neg
|| true positi

encountere
shown tha
tively to c
wise leth
Similar an
other cases
falsely prec
important t
not only fc

Table XI. Dog as a predictor for organ-specific toxicity in man

Organ system	TP* (%)	FP† (%)	TN‡ (%)	FN§ (%)	FN	No. of compounds
					TP + FN (%)	
Injection site	16	36	40	8	33	25
Integument	12	32	40	16	57	25
Cardiovascular	28	24	36	12	30	25
Respiratory	16	64	16	4	20	25
Bone marrow	80	12	0	8	9	25
Lymphoid	4	72	24	0	0	25
Gastrointestinal	92	8	0	0	0	25
Liver	52	44	4	0	0	25
Renal	32	56	4	8	20	25
Neuromuscular	24	60	12	4	14	25

*True positive, toxicity was observed in both the dogs and in man.

†False positive, toxicity was observed in the dogs but not in man.

‡True negative, no toxicity was observed in the dogs and man.

§False negative, toxicity was not observed in the dogs but was recorded in man.

|| False negatives —corrected false negative, an index of false negative prediction which analyzes for true positives plus false negatives only those compounds which produced the specific toxicity in man.

Table XII. Monkey as a predictor for organ-specific toxicity in man

Organ system	TP* (%)	FP† (%)	TN‡ (%)	FN§ (%)	FN	No. of compounds
					TP + FN (%)	
Injection site	13	26	52	9	40	23
Integument	13	17	57	13	50	23
Cardiovascular	22	26	30	22	50	23
Respiratory	13	48	30	9	40	23
Bone marrow	83	13	0	4	5	23
Lymphoid	0	31	65	4	100	23
Gastrointestinal	74	9	0	17	19	23
Liver	52	35	13	0	0	23
Renal	35	48	13	4	11	23
Neuromuscular	22	30	39	9	28	23

*True positive, toxicity was observed in both the monkeys and in man.

†False positive, toxicity was observed in the monkeys but not in man.

‡True negative, no toxicity was observed in the monkeys and man.

§False negative, toxicity was not observed in the monkeys but was recorded in man.

|| False negatives —corrected false negative, an index of false negative prediction which analyzes for true positives plus false negatives only those compounds which produced the specific toxicity in man.

encountered.¹⁰⁴ In addition, it has been shown that penicillin can be used effectively to cure the guinea pig of an otherwise lethal infection of leptospirosis.⁴⁴ Similar analysis can be made of many other cases in which animals seemingly falsely predicted for toxicity. It becomes important to re-examine the animal data not only for the study of these negative

predictive aspects but also for documentation of valuable positive contributions.

Although the literature contains a number of thorough reviews and discussions on the subject of animal prediction of toxicity, a considerable "data gap" exists. Three previous comparative studies have been reported and are worthy of review. Litchfield^{61, 62} reported on his analysis of

ies that failed to predict
thrombocytopenia in man

y

y

nonkey

y

y (not measured in dog)

y

y

nonkey

nonkey

nonkey

many years. The data is not a direct drug test in part, related to the normal gram-positive flora with an overgrowth with gram-negative, the guinea pig died for one of the principles associated with the man—superinfection; i.e., overgrowth with *us* is more commonly

Table XIII. The combination of dog and monkey as a predictor for organ-specific toxicity in man

Organ system	TP* (%)	FP† (%)	TN‡ (%)	FN§ (%)	FN	No. of compounds
					TP + FN (%)	
Injection site	16	36	40	8	33	25
Integument	24	36	36	4	14	25
Cardiovascular	36	32	28	4	10	25
Respiratory	16	76	4	4	20	25
Bone marrow	88	12	0	0	0	25
Lymphoid	4	76	20	0	0	25
Gastrointestinal	92	8	0	0	0	25
Liver	52	48	0	0	0	25
Renal	36	56	4	4	10	25
Neuromuscular	24	60	12	4	14	25

*True positive, toxicity was observed in both the animals and in man.

†False positive, toxicity was observed in the animals but not in man.

‡True negative, no toxicity was observed in the animals and man.

§False negative, toxicity was not observed in the animals but was recorded in man.

|| False negatives —corrected false negative, an index of false negative prediction which analyzes for true positives plus false negatives only those compounds which produced the specific toxicity in man.

six unnamed drugs which included "an antibiotic, a synthetic antibacterial agent, a tranquilizer, a central nervous system depressant, a chemical which blocks oxidation of alcohol, and a glucocorticoid." The compounds were fed to dogs for at least six months and to rats for one year, and, though doses are not stated, the reader is assured that "dosage levels were used which exceeded the tolerance of the animals." Information on the response in man consisted of 500 or more case reports for each compound. Analysis was made on 39 equally weighted physical signs or hematologic parameters, giving a total of 234 possible signs for the six drugs. Analysis of the data demonstrates that the rat predicted for 34 per cent of the 53 total toxic signs observed in man, thus leaving a 66 per cent corrected false negative index. The dog predicted for 55 per cent of total positive signs, while the composite of dog and rat data predicted for only 57 per cent of the cumulative positive signs with 43 per cent corrected false negatives. When the criteria for predictive toxicity were limited to a sign seen in both dog and rat, the result was the lowest forecast

for the total number of toxic signs in man, 32 per cent, leaving a significant 68 per cent corrected false negative correlation. This study further analyzes this latter group for incidence of correct prediction as a per cent of the total positive predictions made. It is stated that toxicity in dog and rat as a predictor showed "... 68 per cent of the positive prediction and 79 per cent of the negative prediction were right, for an over-all score of 74 per cent."⁶² This was interpreted as demonstrating that a toxic parameter shown in two species would more likely predict for toxicity in a third. However, this type of analysis distorts the data and gives a false impression of high correlation with man. It represents only the efficiency of a positive prediction, the number of times it predicted correctly of those times it predicted for toxicity, but fails to take into account the number of false negatives, which in this case represents 26 signs or 68 per cent of the toxicity seen in man.

Of great interest was the remarkable duplication in the rat and dog data. Eighteen out of 234 toxic signs were seen in the rat, while 29 were detected in the

an-specific

FN	No. of compounds
TP + FN (%)	
33	25
14	25
10	25
20	25
0	25
0	25
0	25
0	25
10	25
14	25

prediction which analyzes for
city in man.

of toxic signs in man,
g a significant 68 per
negative correlation.
analyzes this latter
of correct prediction
total positive predic-
ed that toxicity in dog
tor showed ". . . 68
tive prediction and 79
gative prediction were
all score of 74 per
interpreted as demon-
parameter shown in
more likely predict for
However, this type of
data and gives a false
correlation with man.
ie efficiency of a posi-
number of times it
of those times it pre-
but fails to take into
er of false negatives,
represents 26 signs or
icity seen in man.

was the remarkable
it and dog data. Eigh-
ic signs were seen in
were detected in the

Table XIV. The use of single liver function parameters in dogs and monkeys as
a predictor for hepatotoxicity in man

	TP*	FP†	TN‡	FN§	FN TP + FN
<i>Dog</i>					
BSP ↑	5/18¶ 27.8%	9/18 50.0%	1/18 5.6%	3/18 16.7%	3/8 37.0%
Alkaline phosphatase ↑	8/24 33.3%	10/24 41.7%	4/24 16.7%	2/24 8.3%	2/10 20.0%
SGOT ↑	5/18 27.8%	7/18 38.9%	4/18 22.2%	2/18 11.1%	2/7 28.0%
SGPT ↑	3/11 27.3%	5/11 45.5%	3/11 27.3%	0/11	0/3
Bilirubin ↑	3/16 18.8%	3/16 18.8%	7/16 43.8%	3/16 18.8%	3/6 50.0%
Prothrombin time ↑	3/21 14.3%	1/21 4.8%	10/21 47.6%	7/21 33.3%	7/10 70.0%
<i>Monkey</i>					
BSP ↑	3/9 33.3%	4/9 44.4%	1/9 11.1%	1/9 11.1%	1/4 25.0%
Alkaline phosphatase ↑	3/20 15.0%	2/20 10.0%	8/20 40.0%	7/20 35.0%	7/10 70.0%
SGOT ↑	6/18 33.3%	4/18 22.2%	6/18 33.3%	2/18 11.1%	2/8 25.0%
SGPT ↑	3/13 23.1%	3/13 23.1%	6/13 46.2%	1/13 7.7%	1/4 25.0%
Bilirubin ↑	0/14	2/14 14.3%	6/14 42.9%	6/14 42.9%	6/6 100.0%
Prothrombin time ↑	0/6	1/6 16.7%	3/6 50.3%	2/6 33.3%	2/2 100.0%
<i>Dog and monkey</i>					
BSP ↑	5/19 26.3%	9/19 47.4%	1/19 5.3%	4/19 21.1%	4/9 44.0%
Alkaline phosphatase ↑	8/25 32.0%	10/25 40.0%	4/25 16.0%	3/25 12.0%	3/11 28.0%
SGOT ↑	7/20 35.0%	8/20 40.0%	4/20 20.0%	1/20 5.0%	1/8 12.5%
SGPT ↑	4/13 30.1%	6/13 46.2%	3/13 23.1%	0/13	0/4
Bilirubin ↑	3/18 16.7%	4/18 22.2%	7/18 30.9%	4/18 22.2%	4/7 57.0%
Prothrombin time ↑	3/21 14.3%	2/21 9.5%	9/21 42.8%	7/21 33.3%	7/10 70.0%

*True positive, toxicity was observed in both the animals and in man.

†False positive, toxicity was observed in the animals but not in man.

‡True negative, no toxicity was observed in the animals and man.

§False negative, toxicity was not observed in the animals but was recorded in man.

|| False negatives

|| true positives plus false negatives —corrected false negative, an index of false negative prediction which analyzes for

only those compounds which produced the specific toxicity in man.

¶ Numerator = number of drugs producing toxicity

denominator = number of drugs tested

Table XV. Reaction of the alkaline phosphatase and serum transaminase determinations when BSP retention gave a true positive correlation for liver toxicity

	Acetophenone, 2-(dimethyl- amino)-3' 4'-dihydroxy-, hydrochloride	Pactamycin	9H-Purine, 6- (methylthio)-9- β -D-ribofura- nosyl-, dihy- drate	Tubercidin	Urea, 1,3-bis (2-chloroethyl)- 1-nitroso-
<i>Dog</i>					
Alkaline phosphatase	TP	TP	TP	TP	TP
SGOT	TP	FN	—	—	TP
SGPT	—	TP	—	—	TP
<i>Monkey</i>					
Alkaline phosphatase	FN	FN	FN	TP	FN
SGOT	TP	FN	TP	—	TP
SGPT	—	FN	TP	—	—
<i>Dog and monkey</i>					
Alkaline phosphatase	TP	TP	TP	TP	TP
SGOT	TP	FN	TP	—	TP
SGPT	—	TP	TP	—	—

TP = true positive; FN = false negative; — = test not performed.

Table XVI. Reaction of the alkaline phosphatase and serum transaminase determination when BSP retention gave a false negative correlation for liver toxicity

	Cytosine, 1- β -D- arabinofuranosyl-, monohydro- chloride	Hydrazine, 1-acetyl- 2-picolinoyl-	Imidazole-4(or 5)- carboxamide, 5(or 4)-(3, 3-dimethyl-1- triazeno)-	Tylocerebrine
<i>Dog</i>				
Alkaline phosphatase	TP	—	TP	FN
SGOT	—	TP	TP	FN
SGPT	—	TP	—	—
<i>Monkey</i>				
Alkaline phosphatase	FN	FN	TP	FN
SGOT	—	TP	FN	TP
SGPT	—	TP	—	—
<i>Dog and monkey</i>				
Alkaline phosphatase	TP	FN	TP	FN
SGOT	—	TP	TP	TP
SGPT	—	TP	—	—

TP = true positive; FN = false negative; — = test not performed.

dog. When the rat and dog data were combined, the total number of signs observed was 30. Thus, the rat added very little and compared with the dog gave a significantly lower percentage of true

positives and a higher incidence of false negatives for toxicity in man. The list of toxicities unpredicted by the animal screen included, in addition to symptoms, such parameters as diarrhea, gastrointesti-

nal infla
bocytop

This :
analysis
eral de
were us
to inclu
function:
actual
ployed,
duced in
animal s
cult. The
cal parar
used for
stated. In
paramete
possible
an organ
abnormal
rameter.

Owens,
view, and
for bone
renal, neu
compound
of varied
fication, a
ployed for
system pre
cluded mi
data on m
information
therapy, re
of animals,
ria for tox
both roder
torily for g
but failed
The dog pi
testinal and
ing definite
the latter c
gave good
but neither
tory indica
skin and hai

Extending
Pinkel,⁸⁴ Fr
ducted a re

ninase
liver

	Urea, 1,3-bis (2-chloroethyl)- 1-nitroso-
mercaptopurine	
TP	TP
—	TP
—	TP
TP	FN
—	TP
—	—
TP	TP
—	TP
—	—

ase
liver

for 5)- ide, '3, il-1-)-	Tylocrebrine
	FN
	FN
	—
	FN
	TP
	—
	FN
	TP
	—

er incidence of false
ty in man. The list
icted by the animal
addition to symptoms,
diarrhea, gastrointesti-

nal inflammation, aplastic anemia, throm-
bocytopenic purpura, and dermatitis.

This study, while valuable as a pioneer
analysis of animal predictability, has sev-
eral deficiencies. Only six compounds
were used, though there was an attempt
to include a wide range of chemical and
functional types. The lack of details on the
actual compounds, dosage range em-
ployed, maximum degree of toxicity pro-
duced in the animals, and size of the
animal sample make interpretation diffi-
cult. The clinical hematologic and chemi-
cal parameters monitored and the criteria
used for designation of toxicity were not
stated. In addition, analysis by individual
parameters alone made no allowance for
possible correct prediction of toxicity in
an organ, which may be indicated by an
abnormality in a different specific pa-
rameter.

Owens,⁷⁹ in his species comparison re-
view, analyzed the animal predictability
for bone marrow, gastrointestinal, hepatic,
renal, neurologic, and skin toxicities. The
compounds studied were anticancer drugs
of varied chemical and functional classi-
fication, and three to thirteen were em-
ployed for the estimation of each organ
system prediction. The animal species in-
cluded mice, rats, and dogs, with limited
data on monkeys. There was no specific
information given on doses, duration of
therapy, route of administration, number
of animals, parameters followed, or crite-
ria for toxicity. The analysis showed that
both rodent and dog predicted satisfactorily
for general bone marrow depression but
failed to detect thrombocytopenia. The
dog predicted well for both gastroin-
testinal and hepatic toxicity, demonstrat-
ing definite superiority over the rodent in
the latter category. Both rodent and dog
gave good correlation for renal toxicity,
but neither species proved to be satisfac-
tory indicators for either neurologic or
skin and hair toxicity.

Extending the prior observations of
Pinkel,⁸⁴ Freireich and associates⁸⁵ con-
ducted a retrospective study designed to

demonstrate the ability of certain mam-
malian species to predict the toxic dose
levels of anticancer drugs for man. The
analysis of the data from eighteen com-
pounds revealed that the ratio of the
animal-human dose for comparable quan-
titative toxicity approached unity when
measured on the basis of average body
surface area for the species (milligrams
per square meter). A set of factors was
proposed by which a dose in milligrams
per kilogram for one species could be
used to closely estimate a dose which
would produce a similar degree of toxicity
in another animal species or in man. For
example, the maximum tolerated dose
(MTD) in man (milligrams per kilo-
gram) approximates $\frac{1}{12}$ the LD₅₀ in mice,
 $\frac{1}{9}$ the LD₅₀ in hamsters, $\frac{1}{7}$ the LD₅₀ in
rats, $\frac{1}{3}$ the MTD in rhesus monkeys, and
 $\frac{1}{2}$ the MTD in dogs. The study supported
the concept that animal systems can be
effectively used to evaluate the quantita-
tive toxicity of anticancer drugs prior to
use in man.

The present study of qualitative toxicity
deals with twenty-five anticancer drugs of
diverse chemical structure and biochemi-
cal function which have been developed
and/or evaluated by the Chemotherapy
Program of the National Cancer Institute.
The toxicologic evaluation of new com-
pounds in animals has been an essential
facet of this program for several reasons.
A significant proportion of the patients in
whom these compounds were tested were
severely debilitated from disease and thus
rendered less tolerant to drug toxicity. In
this situation perhaps more than in any
other general disease category, the clini-
cian must be alerted to all potential
hazards to his patients. The problem is
made complex by the fact that cancer
chemotherapy drugs, to varying degrees,
fail to discriminate efficiently between
normal and target tissues. A wide range
of toxicities is encountered, and with
many compounds administration of toxic
doses may be required in order to achieve
a therapeutic response. This situation pro-

vided a unique opportunity for the study of drug toxicity per se and facilitated the evaluation of predictions made from animal data.

It is important to note the problems that were encountered in the formulation and analysis of this study. The animal data were collected in both mongrel and beagle dogs and four species of primates. While there appeared to be no differences in predictions in these subgroups, this could not be quantitatively ascertained because of the relatively small numbers in each individual animal group. Normal healthy animals were used in the toxicology studies, while the patients, as a group, were generally debilitated and some measurements were abnormal prior to therapy. In addition, many of the patients were receiving concurrent analgesics, hypnotics, and antiemetics. The animal studies were generally carried out along the format of the CCNSC protocol,¹⁵ which involved the delineation of acute and subacute qualitative toxicities. The animals were generally not treated for longer than 28 days, but this is longer than the average duration of therapy in the clinical studies in man. The route of administration in both animal and man was comparable, but the schedule and dose on both a milligrams per kilogram and milligrams per square meter basis varied (Tables III and IV). The animal and human data were collected by many different laboratories and institutions. While the clinical observations, laboratory determinations, and pathology readings were all considered to be of high quality, they cannot be considered uniform. A set of criteria for estimation of toxicity was formulated, but flexible judgment was employed on the final decision, depending on the animal's pretreatment hematologic and blood chemistry values. No attempt was made to adjust for the incidence of a given specific toxicity observed with each individual compound. It is important that serious toxicity be predicted even though it is subsequently found that its relative incidence in clinical

studies is low. The estimation and comparison of incidence are made impractical, if not impossible, by several factors. The total number of animals receiving the drug was generally significantly lower than the total patient population tested. Treatment in the clinical studies was carried out until the first dose-limiting toxicity was encountered. At this time therapy was halted with the specific intent to prevent further adverse drug reaction. This is in marked contrast to the animal studies in which very toxic and lethal dose ranges had to be administered in order to complete the spectrum of qualitative toxicities inherent with any compound. An additional factor that complicates the usefulness of incidence measurements is that the order of appearance of the qualitative toxicities observed in the different species, as a function of increasing dose, may differ markedly. Such was the case with BCNU, 1,3-bis(2-chloroethyl)-1-nitrosourea, in which the first dose-limiting toxicity noted in dogs was liver damage followed by bone marrow depression. In patients who inadvertently sustained severe drug toxicity the sequence was the reverse, with leukopenia and thrombocytopenia being the predominant toxicities at or near the maximum tolerated dose. Thus, a complete analysis of incidence could be obtained only by carrying out clinical studies in a manner analogous to that of the animal experiment.

We have attempted to establish some of the limitations and useful properties of the large animal toxicology screen by analyzing for prediction of both individual parameters and for organ system toxicity. The species were studied individually and in combination in order to determine whether there was an additive overlap in prediction. This would be represented by an over-all reduction in false negative prediction made by the individual animal species. Such a complementary effect was noted for nine of the ten organ system groups. Perhaps most important was the case of bone marrow depression. The dog

had ur
the mc
suppres
binning
negativ
favorab
for the
The de
leukope
Two of
were co
neither
cator of
demonst
the mor
nine of
depressi
still serv
five fals
all addit
the pred
trast to
the rat in

The o
showed :
in the i
icity, in
cent unc
to none
shown to
productio
observed
twenty-five
demonstra
rhea, cou
negative
dog and
an over-all
relation f
man, this
was contril

Liver t
dicted, bu
quency of
source of
termination
the useful
cedure, eac
mals was
tion of live

stimulation and come made impractical, several factors. The animals receiving the significantly lower population tested. The clinical studies was car-dose-limiting toxic- at this time therapy specific intent to pre- drug reaction. This is the animal studies d lethal dose ranges ered in order to um of qualitative any compound. An complicates the use- measurements is that ce of the qualitative he different species, using dose, may dif- was the case with (methyl)-1-nitrosourea, ose-limiting toxicity er damage followed ession. In patients stained severe drug ; was the reverse, l thrombocytopenia : toxicities at or near l dose. Thus, a com- dence could be ob- g out clinical studies s to that of the ani-

to establish some of ful properties of the y screen by analyz- of both individual rgan system toxicity. lied individually and order to determine additive overlap in d be represented by i in false negative he individual animal lementary effect was e ten organ system : important was the depression. The dog

had underpredicted for two drugs, while the monkey failed to indicate the myelo-suppressive properties of a third. By combining the dog and monkey data, all false negative predictions were removed. This favorable overlap was also demonstrated for the individual hematologic parameters. The dog had failed to predict for the leukopenia produced by five compounds. Two of these false negative predictions were corrected by the monkey. Though neither species proved an adequate indi-cator of thrombocytopenia, the dog was demonstrated to be the superior predictor, the monkey having failed to predict for nine of sixteen drugs producing platelet depression in man. However, the monkey still served to correct for two of the dog's five false negative predictions. The over-all additive effect of the monkey data to the predictions made by the dog is in contrast to the minor contribution made by the rat in Litchfield's studies.

The other category in which the dog showed superiority over the monkey was in the indication of gastrointestinal tox-icity, in which the monkey had a 17 per cent underpredictive index as compared to none for the dog. The monkey was shown to be remarkably resistant to the production of vomiting, the sign that was observed in man with 84 per cent of the twenty-five compounds. The monkey also demonstrated a lower incidence of diar-rhea, coupled with a 26 per cent false negative prediction. While the combined dog and monkey toxicologic screen gave an over-all 92 per cent true positive correlation for gastrointestinal toxicity in man, this high degree of correspondence was contributed by the dog data alone.

Liver toxicity was never underpre-dicted, but there was a 48 per cent fre-quency of false positives. An important source of overprediction was the BSP determination. As an initial investigation of the usefulness of this time-consuming pro-cedure, each liver function test in the ani-mals was analyzed for its over-all predic-tion of liver toxicity in man. It was found

that by using both species and the alka-line phosphatase and a serum transami-nase determination, all cases in which the BSP gave a true positive were similarly predicted; while all BSP false negatives were corrected to positivity. Thus, for the compounds studied, the BSP determina-tion served only as a confirmatory test for liver toxicity and failed to augment infor-mation already obtained by using the transaminase and alkaline phosphatase. Because of the small number of drugs in-volved in each case, a definitive statement regarding the usefulness of the BSP de-termination cannot be made. The alkaline phosphatase in the monkey proved to be a particularly poor determinant of liver toxicity. The range of normal within the species and in individual animals tends to be quite broad; with the use of the pres-ent liberal criteria, a significant percent-age of hepatotoxicity in man is not pre-dicted. The data suggest that these tests, like many others currently employed, should undergo quantitative evaluation for efficacy.

While the animal toxicologic screen failed to predict for only one of ten com-pounds that produced renal toxicity in the clinical studies, this was accomplished at the expense of a 56 per cent false positive index. The high percentage of BUN eleva-tion noted in the animal studies may, in part, reflect extrarenal factors such as gas-trointestinal bleeding and dehydration, but a greater susceptibility of the animal kidney to drug effects was borne out by correlative increases in serum creatinine and histopathologic changes. The inci-dence of false prediction as a function of the total number of compounds producing specific organ toxicity in man ranged from 5 per cent for respiratory, 10 per cent for cardiovascular, to 14 per cent for neurologic and dermal-related toxicities.

In the case of neurologic toxicity, sev-eral general factors become apparent dur-ing the analysis of the data. When compared with the entire spectrum of toxicities inherent in a compound, objec-

tive neurologic toxicity required the largest dose levels for elicitation. The form of the toxicity observed generally differed greatly from that seen in man. Signs such as personality change and confusion cannot be directly predicted by the animals but may be represented by the development of stupor, ataxia, or seizures.

Cardiopulmonary toxicity represents a difficult predictive area, primarily because it is not known which parameter or parameters will be chosen by an animal to express a harmful drug effect. The present study demonstrated a heavy reliance on pathologic changes in organs of the animals in order to predict for clinical parameters in man. In an attempt to develop a better model, future protocols should be designed which include both short- and long-term monitoring of a battery of physiologic parameters including blood pressure, cardiac output, electrocardiogram, and pulmonary x-rays.

The combined large animal screen failed to predict for two of the six compounds that produced injection site toxicity in man. This is explained in part by the failure of the animals to indicate in some way that the compound might produce pain or burning with injection. This symptom was a major problem with the administration of streptozotocin in patients and required that the drug be diluted and infused. The dogs and monkeys showed apparent complete tolerance to preparations of comparable pH (4.5) and buffer molarity.

The animal screen failed to predict dermatitis or correlate well with dermal manifestations of systemic toxicity observed in patients. This deficiency in prediction has long plagued toxicologists. Fortunately, dermatitis, though an unpleasant reaction, rarely poses a life-threatening problem as is more consistently encountered with bone marrow depression.

It is clear that toxicologic data collected in a dog and monkey screen forewarn the clinician of a useful proportion of the total spectrum of organ-specific and, with cer-

tain stated limitations, specific parameter toxicities that might be encountered. With the possible exception of central nervous system and dermal toxicity, all serious organ system toxicities are well predicted. This is accomplished, as in the case of anemia and hepatotoxicity, at the expense of a high percentage of false positive predictions. This degree of inefficiency in the system is justifiable. In order to demonstrate all possible qualitative toxicities, the animals must be given a spectrum of doses including severely toxic and lethal doses. This has been shown for the prediction of bone marrow depression, hepatotoxicity, and neurologic toxicity in our studies. If in the clinic the drugs had been given at more toxic dose levels than the estimated maximum tolerated dose, it is quite possible that the frequency of false positive predictions by the animal species would be significantly lower. Some of the overprediction is likely the result of a more extensive toxicologic evaluation in the animals. It is not common practice in clinical trials to perform lymph node and pulmonary biopsies for histologic evaluation during or immediately after chemotherapy. Both of these parameters showed a high frequency of pathology in the animals but were rarely recorded in man.

In addition to large toxic doses, chronic low-dose administration to animals will serve to elicit toxicities that require repeated subacute insult. These drug effects might be overlooked if an animal is prematurely killed by an earlier appearing, severe qualitative toxicity. An extended observation period of at least two months is required in order to predict for the unusual compound whose toxicity is delayed in onset. Such was the case with BCNU, 1,3-bis(2-chloroethyl)-1-nitrosourea, in the present study. Once organ system toxicity is observed during an animal study, it can never be disregarded but should be viewed with a realistic understanding of certain limitations of animal toxicologic data: Toxicity may develop in man in an organ system predicted susceptible by the animal, but it may be

ons, specific parameter be encountered. With ion of central nervous toxicity, all serious ies are well predicted. ed, as in the case of oxicity, at the expense e of false positive pre e of inefficiency in the . In order to demon- ualitative toxicities, the en a spectrum of doses toxic and lethal doses. n for the prediction of ession, hepatotoxicity, city in our studies. If gs had been given at els than the estimated dose, it is quite possi- ency of false positive animal species would er. Some of the over- the result of a more evaluation in the an- ion practice in clinical mph node and pulmo- tologic evaluation dur- after chemotherapy. meters showed a high ogy in the animals but in man.

ge toxic doses, chronic tion to animals will ities that require re- ult. These drug effects d if an animal is pre- an earlier appearing, toxicity. An extended of at least two months r to predict for the whose toxicity is de- h was the case with chloroethyl)-1-nitroso- it study. Once organ erved during an ani- never be disregarded ed with a realistic un- n limitations of animal xicity may develop in system predicted sus- imal, but it may be

expressed in a different specific clinical or chemical parameter. The adverse reaction may appear in man at a greater or lesser dose level or may follow a different order of appearance in relationship to the total spectrum of qualitative toxicity inherent in any compound. The effective use of animal toxicologic data, coupled with careful monitoring, considered judgment, and expectation by the physician, jointly serve to forewarn the development of critical organ system toxicities during an initial Phase I drug trial.

We would like to acknowledge the following investigators whose personal communications of data were used in the preparation of this paper: Dr. J. Boiron, Dr. J. Hartmann, Dr. A. Haut, Dr. J. Holland, Dr. R. O. Johnson, Dr. D. A. Karnofsky, Dr. M. Lane, Dr. J. Laszlo, Dr. J. Luce, Dr. T. J. Medrek, Dr. T. Necheles, Dr. K. Olson, Dr. A. Schilling, Dr. A. Serpick, Dr. J. Steinfeld, Dr. H. S. Taylor, and Dr. S. A. Taylor.

References

1. Acs, G., Reich, E., and Mori, M.: Biological and biochemical properties of the analogue tubercidin, *Proc. Nat. Acad. Sci.* **52**:493-501, 1964.
2. Ansfield, F. J.: Phase I study of azotomycin (NSC 56654), *Cancer Chemother. Rep.* **46**: 37-40, 1965.
3. Ausman, R. K.: Mitomycin C—Phase II broad spectrum trial, *Proc. Am. A. Cancer Res.* **6**:3, 1965.
4. Barnes, J. M., and Denz, F. A.: Experimental methods used in determining chronic toxicity, *Pharmacol. Rev.* **6**:191-242, 1954.
5. Bhuyan, B. K., Dietz, A., and Smith, C. G.: Pactamycin, new antitumor antibiotic. I. Discovery and biological properties, in *Antimicrobial agents and chemotherapy*, Ann Arbor, 1961, Am. Soc. for Microbiol. pp. 184-190.
6. Bloch, A., and Nichol, C. A.: Studies on the mode of action of 7-deazoadenosine (tubercidin), Abstracts of 145th meeting of the Am. Chem. Soc., Abstract No. P36C, 1963.
7. Bodey, G. P., Brodovsky, H. S., Isassi, A. A., Samuels, M. L., and Freireich, E. J.: Studies of combination 6-mercaptopurine (NSC 755) and 6-methylmercaptopurine riboside (NSC 40774) in patients with acute leukemia and metastatic cancer, *Cancer Chemother. Rep.* **52**:315-320, 1968.
8. Burchenal, J. H.: II. Studies on the mechanism of antileukemic action, in Plattner, P. A., editor: *Present status of the terephthal-*

- anilides, New York, 1964, Elsevier Publishing Company, pp. 233-239.
9. Brodie, B. B.: Distribution and fate of drugs; therapeutic implications, in Binns, T. B., editor: *Absorption and distribution of drugs*, Baltimore, 1964, The Williams and Wilkins Company, pp. 199-251.
10. Brodie, B. B.: Kinetics of absorption, distribution, excretion, and metabolism of drugs, in Nodine, J. H., and Siegler, P. E. editors: *Animal and clinical pharmacologic techniques in drug evaluation*, Chicago, 1964, Year Book Medical Publishers, Inc., pp. 69-88.
11. Brodie, B. B., Cosmides, G. J., and Rall, D. P.: Toxicology and the biomedical sciences, *Science* **148**:1547-1554, 1965.
12. Brodie, B. B., and Reid, W. D.: Some pharmacological consequences of species variation in rates of metabolism, *Fed. Proc.* **26**:1062-1070, 1967.
13. Bryan, G. T., and Gorske, A. L.: Clinical pharmacological studies on hexamethylmelamine (NSC 13875), *Proc. Am. A. Cancer Res.* **9**:9, 1968.
14. Burgen, A. S. V.: The predictive value of animal toxicity tests, *Proc. Second Internat. Pharmacol. Meeting* **8**:49-56, 1965.
15. An outline of procedures for preliminary toxicologic and pharmacologic evaluation of experimental cancer chemotherapeutic agents, *Cancer Chemotherapy National Service Center (CCNSC)*, *Cancer Chemother. Rep.* **37**: 1-33, 1964.
16. Carcy, R. W., and Ellison, R. R.: Continuous cytosine arabinoside infusions in patients with neoplastic disease, *Clin. Res.* **13**:337, 1965.
17. Casazza, A. R., Cahn, E. L., and Carbone, P. P.: Preliminary studies with dibromomannitol (NSC 94100) in patients with chronic myelogenous leukemia, *Cancer Chemother. Rep.* **51**:91-97, 1967.
18. Cassinelli, G., and Orezzi, P.: Daunomycin: A new antibiotic with cytostatic activity, isolation, and properties, *Gior. microbiol.* **11**: 167-174, 1963.
19. Cavins, J. A., Hall, T. C., Olson, K. B., Khung, C. L., Horton, J., Colsky, J., and Shadduck, R. K.: Initial toxicity study of Sangivamycin (NSC 65346), *Cancer Chemother. Rep.* **51**:197-200, 1967.
20. Creasey, W. A., Papac, R. J., Mirkiw, M. E., Calabresi, P., and Welch, A. D.: Biochemical and pharmacological studies with 1- β -D-arabinofuranosylcytosine in man, *Biochem. Pharmacol.* **15**:1417-1428, 1966.
21. Curreri, A. R., and Ansfield, F. J.: Mithramycin—human toxicology and preliminary therapeutic investigation, *Cancer Chemother. Rep.* **8**:18-22, 1960.
22. Curreri, A. R., Ansfield, F. J., Melver, F. A., Waisman, H. A., and Heidelberger, C.: *Clini-*

- cal studies with 5-fluorouracil, *Cancer Res.* 18:478-484, 1958.
23. Dearborn, E. H.: Comparative toxicity of drugs, *Fed. Proc.* 26:1075-1077, 1967.
 24. DeBoer, C., Dietz, A., Lummis, N. E., and Savage, G. M.: Porfiromycin, a new antibiotic. I. Discovery and biological activities, in *Antimicrobial agents annual*, 1960, Washington, D. C., 1961, Am. Soc. for Ind. Microbiol., pp. 17-22.
 25. DeSomer, P., Van De Voorde, H., Eyssen, H., and Van Dijck, P.: A study on penicillin toxicity in guinea pigs, *Antibiotics and Chemother.* 5:463-469, 1955.
 26. DeVita, V. T., Carbone, P. P., Owens, A. H., Jr., Gold, C. L., Krant, M. J., and Edmonson, J.: Clinical trials with 1,3-bis(2-chloroethyl)-nitrosourea, NSC 409962, *Cancer Res.* 25:1876-1881, 1965.
 27. DeVita, V. T., Denham, C., Davidson, J. D., and Oliverio, V. T.: The physiological disposition of the carcinostatic 1,3-bis(2-chloroethyl)-1-nitrosourea (BCNU) in man and animals, *CLIN. PHARMACOL. & THERAP.* 8:566-577, 1967.
 28. DeVita, V. T., Gold, C. L., Owens, A. H., and Miller, J. M.: Preliminary studies with 1,3-bis(2-chloroethyl)-1-nitrosourea (BCNU), *Proc. Am. A. Cancer Res.* 5:15, 1964.
 29. Di Marco, A., Boretti, C., Rusconi, A., and Silvestrini, R.: Metabolic degradation and antineoplastic activity of daunomycin, *Int. Cancer Congr., Abst. 9, Congr. 376*, 1966.
 30. Eckhardt, S., Sellei, C., Horvath, I. P., and Institoris, L.: Effect of 1,6-dideoxy-D-mannitol on chronic granulocytic leukemia, *Cancer Chemother. Rep.* 33:57-61, 1963.
 31. Evans, A. E.: Mitomycin C, *Cancer Chemother. Rep.* 14:1-9, 1961.
 32. Ferguson, D., and Humphrey, E.: Mitomycin C, *Cancer Chemother. Rep.* 8:154, 1960.
 33. Foley, H. T., Shnider, B. I., Gold, C. L., Matias, P. I., Colsky, J., and Miller, S. P.: Phase I studies of porfiromycin (NSC 56410), *Cancer Chemother. Rep.* 51:283-293, 1967.
 34. Frank, W., and Osterberg, A. E.: Mitomycin C (NSC 26980)—an evaluation of the Japanese reports, *Cancer Chemother. Rep.* 9:114-119, 1960.
 35. Freireich, E. J., Gehan, E. A., Rall, D. P., Schmidt, L. H., and Skipper, H. E.: Quantitative comparison of toxicity of anticancer agents in mouse, rat, hamster, dog, monkey, and man, *Cancer Chemother. Rep.* 50:219-244, 1966.
 36. Gale, G. R.: Effect of 1,3-bis(2-chloroethyl)-1-nitrosourea on *Saccharomyces cerevisiae*, *Proc. Soc. Exper. Biol. & Med.* 119:1004-1010, 1965.
 37. Gale, G. R.: Effect of 1,3-bis(2-chloroethyl)-1-nitrosourea on Ehrlich ascites tumor cells, *Biochem. Pharmacol.* 14:1707-1712, 1965.
 38. Glycine, N-diazoacetylhydrazide (NSC 58400), Clinical Brochure, National Cancer Institute, March 12, 1965.
 39. Golberg, L.: The predictive value of animal toxicity studies carried out on new drugs, *J. New Drugs* 3:7-11, 1963.
 40. Gold, C. L., Foley, H. T., and Shnider, B. I.: A preliminary study with porfiromycin (methyl-mitomycin C), *Proc. Am. A. Cancer Res.* 6:22, 1965.
 41. Goldberg, I. H.: Mode of action of antibiotics. II. Drugs affecting nucleic acid and protein synthesis, *Am. J. Med.* 39:722-752, 1965.
 42. Hamre, D. M., Rake, G., McKee, C. M., and MacPhillamy, H. B.: The toxicity of penicillin as prepared for clinical use, *Am. J. M. Sc.* 206:642-652, 1943.
 43. Hananian, J., Holland, J. F., and Sheehe, P.: Intensive chemotherapy of acute lymphocytic leukemia in children, *Proc. Am. A. Cancer Res.* 6:26, 1965.
 44. Heilman, F. R., and Herrell, W. E.: Penicillin in the treatment of experimental leptospirosis icterohemorrhagica, *Proc. Mayo Clin.* 19:89-99, 1944.
 45. Heinemann, B., and Howard, A. J.: Effect of compounds with both antitumor and bacteriophage-inducing activities on *Escherichia coli* nucleic acid synthesis, in *Antimicrobial agents and chemotherapy*, Ann Arbor, 1965, Am. Soc. for Microbiol., pp. 488-492.
 46. Henderson, E. S., and Burke, P. J.: Clinical experience with cytosine arabinoside, *Proc. Am. A. Cancer Res.* 6:26, 1965.
 47. Herr, R. R., Bergy, M. E., Eble, T. E., and Jahnke, H. K.: Porfiromycin, a new antibiotic. II. Isolation and characterization, *Antimicrobial agents annual*, 1960, Washington, D. C., 1961, Am. Soc. for Ind. Microbiol., pp. 23-26.
 48. Hirt, R.: I. Chemical aspects, in Plattner, P. A., editor: *Present status of the terephthalanilides*, New York, 1964, Elsevier Publishing Company, pp. 228-232.
 49. Ho, D. H. W., and Frei, E., III: Comparative biochemical and pharmacological studies of 6-methylthiopurine ribonucleoside (MMPR), *Proc. Am. A. Cancer Res.* 9:31, 1968.
 50. Horwitz, J. P., and Vaitkevicius, V. K.: Purin-6-yltrimethyl-ammonium chloride: A new purine antimetabolite, *Experientia* 17:1-4, 1961.
 51. Howard, J. P., Cevik, N., and Murphy, M. L.: Cytosine arabinoside (NSC 63878) in acute leukemia in children, *Cancer Chemother. Rep.* 50:287-291, 1966.
 52. Institoris, L., Horvath, I. P., Pethes, G., and Eckhardt, S.: Metabolic pathway of cyto-

s
F
53. I
J
s
l
C
54. I
a
a
l
55. Jc
so
E.
ca
2C
Re
56. Jo
rej
an
th
57. Ka
Ce
Re
58. Ki
1-
ic
cel
59. Ko
R.
eml
60. Ko
diff
dru
TH
61. Lit
in
J. A
62. Lit
of r
CLN
196
63. Lit
vent
York
64. Lom
dihy
65. Loo,
W.:
binos
1965
66. Loo,
toxic
Proc.
67. Loo,
J. H
colog
imida
Am.
68. Luce,
Isassi,

- ol. 14:1707-1712, 1965.
- oacetylhydrazide (NSC brochure, National Cancer , 1965.
- predictive value of animal rried out on new drugs, I, 1963.
- y, H. T., and Shnider, B. study with porfiromycin C), Proc. Am. A. Cancer
- Mode of action of antibi- affecting nucleic acid and Am. J. Med. 39:722-752,
- ke, G., McKee, C. M., and B.: The toxicity of peni- for clinical use, Am. J. M. 343.
- ind, J. F., and Sheeche, P.: erapy of acute lympho- children, Proc. Am. A. 1965.
- nd Herrell, W. E.: Peni- ent of experimental lepto- rhagica, Proc. Mayo Clin.
- nd Howard, A. J.: Effect a both antitumor and bac- ng activities on *Escheri-* acid synthesis, in Antimi- l chemotherapy, Ann Ar- c. for Microbiol, pp. 488-
- and Burke, P. J.: Clinical ytosine arabinoside, Proc. . 6:26, 1965.
- , M. E., Eble, T. E., and orfiromycin, a new anti- and characterization, An- nual, 1960, Washington, Soc. for Ind. Microbiol.,
- nical aspects, in Plattner, nt status of the terephthal- , 1964, Elsevier Publishing 232.
- d Frei, E., III: Compara- nd pharmacological studies ne ribonucleoside (MMPR), cer Res. 9:31, 1968.
- and Vaitkevicius, V. K.: -ammonium chloride: A etabolite, Experientia 17:
- vik, N., and Murphy, M. inoside (NSC 63878) in children, Cancer Chemo- 91, 1966.
- ath, I. P., Pethes, G., and tabolic pathway of cyto-
- static dibromohexitols, Cancer Chemother. Rep. 51:261-270, 1967.
53. Iriarte, P. V., Hananian, J., and Cortner, J. A.: Central nervous system leukemia and solid tumors of childhood: Treatment with 1,3-bis(2-chloroethyl)-1-nitrosourea (BCNU), Cancer 19:1187-1194, 1966.
54. Iyer, V. N., and Szybalski, W.: Mitomycins and porfiromycin: Chemical mechanism of activation and cross-linking of DNA, Science 145:55-58, 1964.
55. Johnson, R. O., Bisel, H., Andrews, N., Wil- son, W., Rochlin, D., Segaloff, A., Kremenz, E., Aust, J., and Ansfield, F.: Phase I clinical study of 6- α -methylpregn-4-ene-3, 11, 20-trione (NSC 17256), Cancer Chemother. Rep. 50:671-673, 1966.
56. Jones, R., Jr.: Mitomycin C.: A preliminary report of studies of human pharmacology and initial therapeutic trial, Cancer Chemo- ther. Rep. 2:3-7, 1959.
57. Karnofsky, D. A., and Clarkson, B. D.: Cellular effects of anticancer drugs, Ann. Rev. Pharmacol. 3:376-378, 1963.
58. Kim, J. H., and Eidinoff, M. L.: Action of 1- β -D-arabinofuranosylcytosine on the nucle- ic acid metabolism and viability of HeLa cells, Cancer Res. 25:698-702, 1965.
59. Kofman, S., Medrek, T. J., and Alexander, R. W.: Mithramycin in the treatment of embryonal cancer, Cancer 17:938-998, 1964.
60. Koppanyi, T., and Avery, M. A.: Species differences and the clinical trial of new drugs: A review, CLIN. PHARMACOL. & THERAP. 7:250-270, 1966.
61. Litchfield, J. T., Jr.: Forecasting drug effects in man from studies in laboratory animals, J. A. M. A. 177:104-108, 1961.
62. Litchfield, J. T., Jr.: Evaluation of the safety of new drugs by means of tests in animals, CLIN. PHARMACOL. & THERAP. 3:665-672, 1962.
63. Litchfield, J. T., Jr.: Predictability of con- ventional animal toxicity tests, Ann. New York Acad. Sc. 123:268-272, 1965.
64. Lombardino, J. G.: 4,6-diamino-1-alkyl-1,2- dihydro-s-triazenes, J. M. Chem. 6:213, 1963.
65. Loo, R. V., Brennan, M. J., and Talley, R. W.: Clinical pharmacology of cytosine ara- binoside, Proc. Am. A. Cancer Res. 6:41, 1965.
66. Loo, R. V., and Vaughn, C. B.: Human toxicology and pharmacology of porfiromycin, Proc. Am. A. Cancer Res. 7:43, 1966.
67. Loo, T. L., Stasswender, E. A., Jardine, J. H., and Frei, E., III: Clinical pharma- cological studies on 5-(dimethyltriazeno)- imidazole-4-carboxamide (NSC 45388), Proc. Am. A. Cancer Res. 8:42, 1967.
68. Luce, J. K., Frenkel, E. P., Vietti, T. J., Isassi, A. A., Hernandez, K. W., and Howard,
- J. P.: Clinical studies of 6-methylmercap- topurine riboside, Cancer Chemother. Rep. 51:535-546, 1967.
69. MacDonald, C., Wollner, N., Ghavimi, F., and Zweig, J.: Phase I study of imidazole, carboxamide dimethyltriazeno (ICD), Proc. Am. A. Cancer Res. 8:43, 1967.
70. Maddock, C. L., Handler, A. H., Friedman, O. M., Foley, G. E., and Farber, S.: Pri- mary evaluation of alkylating agent cyclo- hexylamine salt of N,N-bis(2-chloroethyl) phosphorodiamidic acid (NSC 69945; OMF- 59) in experimental antitumor assay systems, Cancer Chemother. Rep. 50:629-639, 1966.
71. Manheimer, L. H., and Vital, J.: Mitomycin C in the therapy of far-advanced malignant tumors, Cancer 19:207-212, 1966.
72. Mason, J. H., Wilson, W. L., Ansfield, F. J., Rochlin, D. B., and Grage, T.: Phase I clinical study of 2-(dimethylamino)-3',4'-dihy- droxyacetophenone, hydrochloride (NSC 62512), Cancer Chemother. Rep. 52:297-299, 1968.
73. Miller, E., Sullivan, R. D., and Chryssochoos, T.: The clinical effects of mitomycin C by continuous intravenous administration, Can- cer Chemother. Rep. 21:129-135, 1962.
74. Montgomery, J. A., and Temple, C.: Synth- esis of potential anticancer agents, XXIII. 9-Aminohypoxanthine and related com- pounds, J. Am. Chem. Soc. 82:4592, 1960.
75. Morris, N. R., and Glaser, G. H.: Effects of a pyrimidine analog, 6-azauracil, on rat elec- troencephalogram and maze running ability, Electroencephalography & Clin. Neurophysi- ol. 11:146-150, 1959.
76. Nathanson, L., Hall, T. C., Rutenberg, A., and Shaddock, R. K.: Clinical toxicologie study of cyclohexylamine salt of N,N-bis(2- chloroethyl) phosphorodiamidic acid (NSC 69945; OMF-59), Cancer Chemother. Rep. 51:35-39, 1967.
77. Nies, B. A., Thomas, L. B., and Freireich, E. J.: Meningeal leukemia: A follow-up study, Cancer 18:546-553, 1965.
78. Oettgen, H. F., Clifford, P., and Burchenal, J. H.: Malignant lymphoma involving the jaw in African children: Treatment with 2- chloro-4'-4''-di-2-imidazolin-2-ylterephthalanil- ide dihydrochloride, Cancer Chemother. Rep. 27:45-54, 1963.
79. Owens, A. H., Jr.: Predicting anticancer drug effects in man from laboratory animal studies, J. Chron. Dis. 15:223-228, 1962.
80. Page, L. B., and Culver, P. J.: A syllabus of laboratory examinations in clinical diag- nosis, Cambridge, 1962, Harvard University Press, pp. 343-349.
81. Papac, R., Creasey, W. A., Calabresi, P., and Welch, A. D.: Clinical and pharamco- logical studies with 1- β -arabinofuranosylcy-

- tosine (cytosine arabinoside), *Proc. Am. A. Cancer Res.* 6:50, 1965.
82. Pactamycin, Clinical Brochure, National Cancer Institute, December 15, 1961.
 83. Parker, G. W., Wiltsie, D. S., and Jackson, C. B., Jr.: The clinical evaluation of PA-144 (mithramycin) in solid tumors of adults, *Cancer Chemother. Rep.* 8:23-26, 1960.
 84. Pinkel, D.: The use of body surface area as a criterion on drug dosage in cancer chemotherapy, *Cancer Res.* 18:853-856, 1958.
 85. Pittillo, R. F., Narkates, A. J., and Burns, J.: Microbiological evaluation of 1,3-bis(2-chloroethyl)-1-nitrosourea, *Cancer Res.* 24:1222-1228, 1964.
 86. Rakieten, N., Rakieten, M. L., and Nadkarni, M. V.: Studies on the diabetogenic action of streptozotocin (NSC 37917), *Cancer Chemother. Rep.* 29:91-98, 1963.
 87. Rall, D. P., Ben, M., and McCarthy, D. M.: 1,3-Bis-chloroethyl-1-nitrosourea (BCNU): Toxicity and initial clinical trial, *Proc. Am. A. Cancer Res.* 4:55, 1963.
 88. Rao, K. V., Cullen, W. P., and Sobin, B. A.: A new antibiotic with antitumor properties, *Antibiotics and Chemother.* 12:182-186, 1962.
 89. Reitemeier, R. J., Moertel, C. G., and Hahn, R. G.: 1,3-Bis(2-chloroethyl)-1-nitrosourea (BCNU) therapy in advanced gastrointestinal adenocarcinoma, *Proc. Am. A. Cancer Res.* 7:59, 1966.
 90. Reitemeier, R. J., Moertel, C. G., and Hahn, R. G.: Mitomycin C therapy of advanced gastrointestinal adenocarcinoma. Comparison of short- and long-term treatment schedules, *Proc. Am. A. Cancer Res.* 8:56, 1967.
 91. Report of Eastern clinical drug evaluation program in abstracts of papers, 9th International Cancer Congress, *Cancer Chemother.* 7:533, 1964. (Abst.)
 92. Schein, P. S., Cooney, D. A., and Vernon, M. L.: The use of nicotinamide to modify the toxicity of streptozotocin diabetes without loss of antitumor activity, *Cancer Res.* 27:2324-2332, 1967.
 93. Schepartz, S. A., Wodinsky, I., and Leiter, J.: Phthalanilides—a new group of potential antitumor agents, *Cancer Chemother. Rep.* 19:1-5, 1962.
 94. Schneiderman, M. A.: The clinical excursion into 5-FU, *J. Chron. Dis.* 15:283-295, 1962.
 95. Shealy, Y. F., Montgomery, J. A., and Laster, W. R., Jr.: Antitumor activity of triazenoimidazoles, *Biochem. Pharmacol.* 11:674-675, 1962.
 96. Shnider, B. I., Frei, E., III, Tuohy, J. H., Gorman, J., Freireich, E. J., Brindley, C. O., Jr., and Clements, J.: Clinical studies of 6-azauracil, *Cancer Res.* 20:28-33, 1960.
 97. Spear, P. W.: Clinical trial with mithramycin, *Cancer Chemother. Rep.* 29:109-113, 1963.
 98. Talley, R. W., Vaitkevicius, V. K., Reed, M. L., and Brennan, M. J.: Cytosine arabinoside; human pharmacology and toxicity, *Proc. Am. A. Cancer Res.* 3:366, 1962.
 99. Tan, C., Tasaka, H., Yu, K. P., Murphy, M. L., and Karnofsky, D. A.: Daunomycin, an antitumor antibiotic, in the treatment of neoplastic disease, *Cancer* 20:333-353, 1967.
 100. Theologides, A., Yarbrow, J. W., and Kennedy, B. J.: Daunomycin inhibition of DNA and RNA synthesis in normal and malignant tissues, *Proc. Am. A. Cancer Res.* 8:67, 1967.
 101. Vaitkevicius, V. K., Reed, M. L., Fox, R. L., and Talley, R. W.: Trimethylpurin-6-yl-ammonium chloride (NSC 51095): Acute toxicity in man—a preliminary report, *Cancer Chemother. Rep.* 27:55-61, 1963.
 102. Vaitkevicius, V. K., Brennan, M. J., Beckett, V. L., Kelly, J. E., and Talley, R. W.: Clinical evaluation of cancer chemotherapy with 5-FU, *Cancer* 14:102-131, 1961.
 103. Watne, A. L., Moore, D., and Gorgun, B.: Solid tumor chemotherapy with mitomycin C, *Proc. Am. A. Cancer Res.* 8:71, 1967.
 104. Weinstein, L.: Superinfection: A complication of antimicrobial therapy and prophylaxis, *Am. J. Surg.* 107:704-709, 1964.
 105. Weiss, A. J., Goldman, L., Ramirez, G., and Hill, G. J., II: A Phase II study of azotomycin, *Proc. Am. A. Cancer Res.* 9:76, 1968.
 106. Weissbach, A., and Lisio, A.: Alkylation of nucleic acids by mitomycin C and porfiromycin, *Biochemistry* 4:196-200, 1965.
 107. Welch, A. D., Handschumacher, R. E., and Jaffe, J. J.: Studies on the pharmacology of 6-azauracil, *J. Pharmacol. & Exper. Therap.* 129:262-270, 1960.
 108. Wheeler, G. P., and Bowdon, B. J.: Some effects of 1,3-bis(2-chloroethyl)-1-nitrosourea upon the synthesis of protein and nucleic acids in vivo and in vitro, *Cancer Res.* 25:1770-1778, 1965.
 109. White, F. R.: New agent data summaries: Mitomycin C, *Cancer Chemother. Rep.* 2:20-21, 1959.
 110. White, F. R.: Pactamycin, *Cancer Chemother. Rep.* 24:75-78, 1962.
 111. Williams, R. T.: Comparative patterns of drug metabolism, *Fed. Proc.* 26:1029-1039, 1967.
 112. Wilson, W., Schroeder, J., Bisel, H., Mrazek, R., and Hummel, R.: Phase II study of hexamethylmelamine (NSC 13875), *Cancer* 23:132-136, 1969.
 113. Zbinden, G.: Advances in pharmacology, vol. 2, New York, 1963, Academic Press, Inc., pp. 1-112.